

Conclusion The two irradiation regimens used in HDR utero-vaginal brachytherapy in 3 or 5 fractions give similar oncological results with a comparable late toxicity.

Disclosures HR CTV= high-risk clinical target volume

EQD2= Equivalent dose in 2Gy fractions

#870

RADIOMIC PROFILES IMPROVE PROGNOSTICATION AND REVEAL TARGETS FOR THERAPY IN CERVICAL CANCER

^{1,2}Mari Kyllèsø Halle*, ^{3,4}Erlend Hodneland, ^{5,6}Kari S Wagner-Larsen, ^{7,5}Njål G Lura, ^{8,9}Kristine E Fasmer, ^{4,10}Hege F Berg, ^{7,5}Thomasz Stokowy, ^{8,9}Aashish Srivastava, ^{4,10}David Forse, ^{7,5}Erling A Hoivik, ⁸Kathrine Woie, ⁴Bjørn I Bertelsen, ^{7,5}Camilla Krakstad, ^{8,2}Ingfrid S Haldorsen. ¹Centre for Cancer Biomarkers, Bergen, Norway; ²Department of Obstetrics and Gynecology, Bergen, Norway; ³Department of Clinical Science, Bergen, Norway; ⁴Haukeland University Hospital, Bergen, Norway; ⁵University of Bergen, Bergen, Norway; ⁶Department of Mathematics, Bergen, Norway; ⁷Mohn Medical Imaging and Visualization Centre, Bergen, Norway; ⁸Department of Radiology, Bergen, Norway; ⁹Section of Radiology, Bergen, Norway; ¹⁰Department of Clinical Medicine, Bergen, Norway; ¹¹Section of Radiology, ¹²Department of Clinical Medicine

10.1136/ijgc-2023-ESGO.198

Introduction/Background Pelvic magnetic resonance imaging (MRI) is an important part of primary diagnostic workup in cervical cancer (CC), with MRI-assessed tumour size and pelvic tumour extent routinely guiding treatment decisions. Extraction of MRI-derived radiomic tumour features could improve cancer prognostics and may also reveal novel targets for treatment.

Methodology We manually segmented 3D volumes in 132 primary CC tumours and extracted 293 whole-volume radiomic MRI features. Unsupervised hierarchical clustering yielded three distinct patient clusters (Cluster 1 [n=52]; 2 [n=46]; and 3 [n=34]). Overlapping clinicopathologic, genomic (whole exome sequencing, n=65), transcriptomic (L1000 arrays, n=73) and molecular biomarker (n=84) data were utilized to characterize each cluster.

Results Patients in Cluster 2 and 3 had significantly reduced disease-specific survival (DSS) (hazard rate [HR]: 3.33; p=0.008) compared with patients in Clusters 1, even after adjusting for International Federation of Gynecology and Obstetrics (FIGO) 2018 stage and age (adjusted HR: 2.51; p=0.045). Cluster 3 tumours associate with high stage (p<0.001), large tumours (p<0.001), squamous histology (p=0.015), p53 negative or -overexpressing tumours (p=0.04) and aberrant TP53, -MYC and -MTORC1 signalling. The intermediate-risk Cluster 2 associates with increased and aberrant cell cycle- and Hippo signalling, suggesting CDK2/4 and YAP-TEAD inhibitors as plausible treatment options. The low-risk Cluster 1 associates with increased immune cell signalling.

Conclusion This study links radiomic signatures to distinct genomic profiles that may potentially aid in prognostication and tailoring of treatments and follow-up plans for cervical cancer patients.

Disclosures The authors declare no conflict interests.

#874

LOCALLY ADVANCED CERVICAL CANCER IN YOUNG WOMEN

¹Marwa Aloui, ¹Ines Zemni, ¹Houyem Mansouri, ¹Souha Jaouadi*, ²Nedia Boujelbene, ¹Tarek Ben Dhiab. ¹Surgical oncology department, Salah Azaiez Institute of oncologie, Tunis, Tunisia; ²Pathology department, Salah Azaiez Institute of oncologie, Tunis, Tunisia

10.1136/ijgc-2023-ESGO.199

Introduction/Background The incidence of cervical cancer in young women is increasing. This study aims to analyze the clinicopathological characteristics, treatment, and prognoses of women aged ≤40 years with locally advanced cervical cancer.

Methodology Medical record data of 25 locally advanced cervical cancer patients aged ≤40 years treated at Salah Azaiez Institute of Oncology between January 2010 and December 2020 were reviewed. The overall survival rate was estimated using the Kaplan–Meier method. Prognosis-related risk factors were analyzed using univariate analyses.

Results The median age at diagnosis in our study was 35.8 years ± 3.13. Among 25 patients, 24 (96%) were diagnosed with squamous cell carcinoma and 1 (4%) with adenocarcinoma. In 12 cases (48%), the FIGO stage was ≤ IIB and in 13 cases (52%) FIGO stage was >IIB. The median tumor size at diagnosis was 54.6mm ± 17.37. 24% of patients had multiple sexual partners. Only 15 patients were treated with curative intentions. 3 patients presented locoregional recurrence. No patient presented metastatic recurrence. 7 patients presented progression during treatment. 17 cases of death during follow-up were noticed among 25 cases.

One-year disease-free survival (DFS) was 67.2%. lympho-vascular space involvement (LVSI) (p=0.010) was a prognostic factor that affected disease-free survival. DFS was not significantly different for tumor size (p=0.0179), cell type (p=0.5) histological grade (p=0.45), parametrial involvement (p=0.068), and pelvic lymph node metastases (p=0.155).

One-year overall survival (OS) was 46.7%. Parametrial involvement (p=0.001), FIGO stage (p=0.028), and histological grade were risk factors for poor prognosis for these patients. OS survival was not significantly different for tumor size, LVSI, and pelvic lymph node metastases.

Conclusion Locally advanced cervical cancer in young women presents an overall prognosis worse than in older patients.

Disclosures We have no potential conflict of interest to report

#878

DEFINITIVE RADIOTHERAPY/CHEMORADIOTHERAPY RESULTS IN GERIATRIC PATIENTS WITH CERVICAL CANCER

¹Ilknur Alsan Cetin, ²Hatice Halis, ³Melis Gültekin, ⁴Senem Alanyali, ⁵Sefika Arzu Ergen, ⁶Kamuran Ibis, ⁷Binnaz Sarper, ⁸Zeliha Güzelöz, ⁹Barbaraos Aydın, ¹⁰Hatice Öner, ¹¹Dile Arslan, ¹²Candan Demiröz Abakay, ¹³Enis Özyar, ¹⁴Alpaslan Seraslan, ¹⁵Havva Beyaz, ¹⁶Aysenur Elmali, ⁴Tugce Bozkurt, ¹⁷Seden Küçükçuk, ⁷Nermin Mirzezade, ³Ferah Yıldiz, ⁴Zeynep Özşaran, ⁹Ilknur Bilkay Görken, ¹⁸Evrin Metcalfe. ¹Marmara University Faculty of Medicine Radiation Oncology Department, Istanbul, Turkey; ²Sakarya Training and Research Hospital, Sakarya, Turkey; ³Hacettepe University Faculty of Medicine Radiation Oncology Department, Ankara, Turkey; ⁴Ege University Faculty of Medicine Radiation Oncology Department, Izmir, Turkey; ⁵Istanbul University Cerrahpasa Medical Faculty Hospital Radiation Oncology Department, Istanbul, Turkey; ⁶Istanbul University Medical Faculty Hospital Radiation Oncology Department, Istanbul, Türkiye; ⁷Kocaeli University Medical Faculty Hospital Radiation Oncology Department, Kocaeli, Turkey; ⁸Tepecik Training and Research Hospital, Department of Radiation Oncology, Izmir, Turkey; ⁹Dokuz Eylul University Faculty of Medicine Radiation Oncology Department, Izmir, Turkey; ¹⁰Aydın Atatürk State Hospital, Department of Radiation Oncology, Aydın, Turkey; ¹¹Erciyes University Faculty of Medicine Radiation Oncology Department, Kayseri, Turkey; ¹²Bursa Uludağ University Hospital Radiation Oncology Department, Bursa, Turkey; ¹³Acibadem University Hospital Radiation Oncology Department, Istanbul, Turkey; ¹⁴Ondokuz Mayıs University Faculty of Medicine Radiation Oncology Department, Samsun, Turkey; ¹⁵Ankara City Hospital Radiation Oncology Department, Ankara, Turkey; ¹⁶Ministry of Health Elazığ Fethi Sekin City Hospital, Elazığ, Turkey; ¹⁷Istanbul University Medical Faculty Hospital Radiation Oncology Department, Istanbul, Turkey; ¹⁸Medipol University Medical Faculty Hospital Radiation Oncology Department, Istanbul, Turkey

10.1136/ijgc-2023-ESGO.200

Introduction/Background To investigate the prognostic factors for survival, the long-term results and toxicities in elderly (≥ 65 years) patients with cervical cancer who underwent definitive radiotherapy/chemoradiotherapy with a cervical cancer diagnosis.

Methodology We reviewed the clinical records of 458 patients with cervical cancer. The median age was 71 (65–91). The median treatment dose was 50.4 Gy (45–64). Overall survival (OS), cancer-specific survival (CSS), disease-free survival (DFS), distant metastasis-free survival (DMFS), and local recurrence-free survival (LRFS) were evaluated using Kaplan-Meier, log-rank, and Cox regression analyses; $p < 0.05$ was considered statistically significant.

Results Median follow-up 48 months (6–243); The 5-year OS, CSS, DFS, DMFS, and LRFS were 61.6%, 77.5%, 75.6%, 81.1%, and 91.6%, respectively. OS, CSS, DFS, DMFS influencing factors performance status (PS) ($p < 0.0001$) in univariate analysis, tumor diameter (≤ 4 cm, > 4 cm) ($p = 0.004$, $p = 0.008$, $p < 0.0001$) histopathology (squamous, adeno and others) ($p = 0.001$, $p < 0.0001$), presence of lymph nodes (LN) ($p < 0.0001$), treatment response (complete, partial, unresponsive) ($p < 0.0001$) was statistically significant. For OS, age (65–70, 71–79, ≥ 80) ($p < 0.0001$), type of treatment (CCRT, RT) ($p < 0.0001$), Brachytherapy (BT) application ($p < 0.0001$); BT application for CSS and DMFS ($p = 0.025$, $p = 0.003$) and treatment response for LRFS ($p < 0.0001$) were significant parameters. In multivariate analysis, independent prognostic factors for OS, CSS, DMFS, and DFS, are treatment response ($p < 0.0001$, $p = 0.003$), LN ($p < 0.0001$, $p = 0.007$, $p = 0.002$, $p = 0.003$) and PS for OS, DFS ($p = 0.003$, $p = 0.004$), the age for OS ($p = 0.045$), type of treatment ($p < 0.0001$). For DMFS, histopathology ($p = 0.026$) and BT application ($p = 0.049$) were found to be significant.

Conclusion Curative treatments for geriatric patients, especially in locally advanced cervical cancer, have a low rate of side effects but have a favorable effect on survival and definite CRT should be recommended in geriatric age.

Disclosures Curative treatments for geriatric patients, especially in locally advanced cervical cancer, have a low rate of side effects but have a favorable effect on survival and definite CRT should be recommended in geriatric age.

#879

SURGERY AFTER CONCURRENT CHEMORADIO THERAPY AND BRACHYTHERAPY IN LOCALLY ADVANCED CERVICAL CANCER: MORBIDITY AND OUTCOME

Marwa Aloui, Ines Zemni, Saida Sakhr, Souha Jaouadi*, Riadh Chargui, Tarek Ben Dhied. *Surgical Oncology Department, Salah Azaiez Institute of Oncologie, Tunis, Tunisia*

10.1136/ijgc-2023-ESGO.201

Introduction/Background This study was performed to evaluate the morbidity and therapeutic value of surgery after concurrent chemoradiotherapy and brachytherapy in patients with locally advanced cervical cancer (LACC).

Methodology We conducted a retrospective study at Salah Azaiez Institute of Oncology from January 1, 2010, to December 31, 2020, including 118 patients with LACC treated with concurrent chemoradiotherapy and pelvic radiotherapy followed by brachytherapy. The surgical treatment consisted of a hysterectomy, which ranged from radical

hysterectomy to anterior pelvic exenteration, and lymph node resection.

Results 118 patients were enrolled over 10 years. The median age was 53 years ± 10.758 . Patients' distribution according to FIGO stage was as follows: 72 (61%) patients $\leq$ IIB and 46 (39%) $>$ IIB. Among these patients, 49 (41.5%) experienced postoperative complications. 9 patients experienced Grade II morbidity while 18 patients presented Grade III morbidity. No postoperative mortality was observed. The median operative time was 215.59 min ± 68.7 . The median duration of hospitalization was 6.12 days ± 15.7 . 41 patients (34.7%) needed per-operative transfusions. The median number of packed red blood cells was 2.46 ± 1.12 . 39 patients presented preoperative complications. Per-operative bleeding was the most common complication observed in 38 cases.

Personal history of High blood pressure ($p = 0.0149$), tumor size ($p = 0.051$), and pelvic anterior exenteration ($p = 0.037$) were risk factors of postoperative morbidity.

Post-operative morbidity was not significantly different for age ($p = 0.781$), FIGO stage ($p = 0.969$), and per-operative transfusion ($p = 0.799$).

Conclusion Surgery after concurrent chemoradiotherapy and brachytherapy for advanced cervical cancer lead to an acceptable morbidity. In our study, Personal history of High blood pressure, tumor size, and pelvic anterior exenteration were risk factors for postoperative morbidity.

Disclosures We have no potential conflict of interest to report.

#886

SUCCESSFUL TOPICAL IMMUNOTHERAPY WITH 5% IMIQUIMOD IN CERVICAL STUMP PREMALIGNANT LESION – A CASE REPORT

¹Damir Danolic, ¹Melita Peric Balja*, ¹Lucija Šušnjari, ¹Ilija Alvir, ¹Ivica Mamic, ²Tomislav Becejac, ³Danijela Danolic, ¹Mario Puljiz. ¹University Hospital for Tumours, University Hospital Centre Sestre milosrdnice, Zagreb, Croatia; ²Department of Thoracic Surgery, University Hospital Centre Zagreb, Zagreb, Croatia; ³School of Dental Medicine, University of Zagreb, Zagreb, Croatia

10.1136/ijgc-2023-ESGO.202

Introduction/Background Imiquimod is an immune response modulator often used in the management of various clinical manifestations of human papillomavirus (HPV) infection. The aim of this case report is to present a possibility of non-surgical, pharmacotherapeutic approach with 5% imiquimod as an appropriate alternative to surgical procedures in selected patients with cervical premalignant lesions.

Results A 50-year-old female presented with cervical stump high grade squamous intraepithelial lesion (HSIL), seven years after laparoscopic supracervical hysterectomy. Current guidelines recommend surgery as preferable treatment modality for histologically confirmed cervical HSIL, after 6 months follow-up Pap test. Initial approach to management of patients with cervical stump HSIL must consider treatment-related morbidity. The goal of treatment is to prevent disease progression to invasive disease. Nonsurgical medical therapy with topical imiquimod appears to be more effective than no treatment, inducing regression of the disease in 73%, versus 50%. Considering these data, after biopsy and during 6 months' period of follow-up, we decided to initiate imiquimod therapy until the follow-up Pap test. Imiquimod treatment was started with self-applied vaginal imiquimod suppositories three times