

# Predictors risk factors for acute complex appendicitis pain in patients: Are there gender differences?

Cem Cahit Barışık<sup>1</sup>, Abdulbari Bener<sup>2,3,4</sup>

Departments of <sup>1</sup>Radiology and <sup>4</sup>Public Health, Medipol School of Medicine, Istanbul Medipol University, <sup>2</sup>Department of Biostatistics and Medical Informatics, Cerrahpaşa Faculty of Medicine, Istanbul University Cerrahpaşa, Istanbul, Turkey, <sup>3</sup>Department of Evidence for Population Health Unit, School of Epidemiology and Health Sciences, The University of Manchester, Manchester, UK

## ABSTRACT

**Objective:** The purpose of this study is to determine the predictive risk factors for appendicitis and the cost-effectiveness of using abdominal helical computed tomography (CT) in comparison to abdominal ultrasonography (US) for the diagnosis of acute appendicitis in patients. **Subjects and Methods:** The typical case was a patient with abdominal pain in the right lower quadrant and suspicion of appendicitis. A total of 643 patients who were consequently treated with appendectomy upon diagnosis of acute appendicitis between January 2015 and December 2018 were included in the study. The four diagnostic alternatives chosen were US, CT, biochemistry parameters, and physical examination in the hospital. **Results:** There were statistically significant differences between male and female patients with regards to age, BMI, cigarette smoking, sheesha smoking, family history of diabetes, hypertension and family history of gastrointestinal discomfort (GI), anxiety ( $P < 0.001$ ), red eye ( $P = 0.006$ ), dizziness ( $P = 0.021$ ), headache ( $P < 0.001$ ), muscular symptoms, weakness and cramps ( $P < 0.001$ ), bloating or swollen stomach ( $P < 0.001$ ), UTI ( $P < 0.001$ ), chest pain ( $P < 0.001$ ), guarding ( $P < 0.001$ ), loss of appetite ( $P = 0.004$ ), nausea ( $P < 0.001$ ) vomiting ( $P = 0.042$ ), anorexia ( $P = 0.009$ ), and constipation ( $P = 0.002$ ). Moreover, there were statistically significant differences between male and female patients for pain ( $P < 0.001$ ), pain right belly ( $P = 0.027$ ), severe cramps ( $P = 0.007$ ), high temperature and fever ( $P < 0.001$ ), irritable bowel syndrome ( $P < 0.001$ ), right iliac fossa (RIF) pain ( $P = 0.008$ ), rebound tenderness ( $P = 0.024$ ), positive bowel sounds ( $P = 0.029$ ), and pointing tenderness ( $P < 0.001$ ). Multivariate stepwise logistic regression showed nausea ( $P < 0.001$ ), C-reactive protein (CRP) ( $P < 0.001$ ), dizziness ( $P = 0.016$ ), vomiting ( $P < 0.001$ ), muscular symptoms ( $P = 0.007$ ), irritable bowel syndrome ( $P = 0.034$ ), guarding ( $P = 0.040$ ), and loss appetite ( $P = 0.046$ ) were considered at higher risk as predictors for appendicitis patients. **Conclusions:** CT is more cost-effective than the US and clinical examination for determining appendicitis. The current study suggested that nausea, C-reactive protein, dizziness, vomiting, muscular symptoms, irritable bowel syndrome, guarding, and loss appetite were considered as higher risk predictors for appendicitis patients.

**Keywords:** Appendicitis, computed tomography, diagnosis, gender, predictors, ultrasound

## Introduction

Acute appendicitis diagnosis is complicated in approximately 35% of patients with pain in the lower right quadrant<sup>[1]</sup> which is the most common cause of surgical abdominal pain.<sup>[2]</sup> Therefore, timely diagnosis of acute appendicitis has an impact on the overall health and economic status of most countries.<sup>[3]</sup> The previously proposed disadvantages of computed tomography (CT) such

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: [WKHLRPMedknow\\_reprints@wolterskluwer.com](mailto:WKHLRPMedknow_reprints@wolterskluwer.com)

**How to cite this article:** Barışık CC, Bener A. Predictors risk factors for acute complex appendicitis pain in patients: Are there gender differences? *J Family Med Prim Care* 2020;9:2688-92.

**Address for correspondence:** Dr. Cem Cahit Barışık, Department of Radiology, İstanbul Medipol University, Medipol Hospital Acıbadem Koşuyolu, Kadıköy, 34718, Istanbul, Turkey. E-mail: [cbarisik@medipol.edu.tr](mailto:cbarisik@medipol.edu.tr)

Received: 21-01-2020

Revised: 13-03-2020

Accepted: 08-04-2020

Published: 30-06-2020

### Access this article online

#### Quick Response Code:



Website:  
[www.jfmpc.com](http://www.jfmpc.com)

DOI:  
10.4103/jfmpc.jfmpc\_140\_20

as it is expensive, not being available everywhere, and the use of contrast medium<sup>[4]</sup> have gradually decreased, and today CT is more commonly used in the diagnosis of appendicitis. Using ultrasonography (US) and CT for assessing acute appendicitis has improved diagnostic accuracy for what can be a difficult clinical diagnosis.<sup>[5-9]</sup> The imaging diagnosis of acute appendicitis can be made accurately by US or CT.<sup>[6-12]</sup> Overall, fortunately, the advances in technology with the development of US and CT have shown considerable advantages in the diagnosis of patients with suspected acute appendicitis.<sup>[13-18]</sup> Several studies reported that both White blood cell (WBC) and C-reactive protein (CRP) proved to be a fair and very poor predictor of complex appendicitis.<sup>[19,20]</sup>

This study aims to determine the predictive risk factors and cost-effectiveness for appendicitis using CT and ultrasound in the diagnosis of acute appendicitis in patients who have consequently been treated with appendectomy upon preliminary diagnosis.

## Subjects and Methods

This prospective cohort study included adult patients between the ages of 20 and 60 who visited the emergency department, gastroenterology, and surgery and outpatient clinics in the İstanbul Medipol University, Faculty of Medicine Teaching Hospitals. The study was conducted between January 2016 and July 2019 using a total of 643 consecutive patients who underwent both CT and appendix the US for suspected acute appendicitis. Institutional Review Board (IRB) ethical approval for the current study was obtained from the Medipol International School of Medicine, İstanbul Medipol University.

### Radiological measurements

#### Ultrasound

A general abdominal examination was performed using sonography. The results of the examination were recorded on a digital case record form; the following potential appendiceal abnormalities on imaging were used as diagnostics for appendicitis: inability visualizing the appendix completely (using General Electric Logic P6 Pro, (transducer) 4 MHz, 5 MHz, and 10 MHz), the presence of local transducer tenderness, the presence of a thickened appendix (diameter greater than 6 mm), and the presence of an incompressible appendix. Most recent study reported<sup>[12]</sup> that the diagnostic performance of ultrasound reevaluation were 96.3% sensitivity, 91.2%, specificity, 89.7% PPV, 96.9%, NPV, and 91.9% accuracy.

#### Computed tomography

CT exams were performed using the General Electric Light speed VCT XT 64 detector helical CT, width 5 mm. The patients based on contrast (nonenhanced) and (enhanced) visualized. CT findings<sup>[12]</sup> provided excellent performance of 96.3% sensitivity, 91.2% specificity, 89.7% PPV, 96.9% NPV, and 91.9% accuracy for diagnosing appendicitis.

The final diagnosis was based altogether on clinical physician examination, laboratory, surgical, pathological histopathology reports, radiological diagnostics with US and CT, and measurements.

The Student's *t*-test was performed for significant differences between the mean of two continuous values and the Chi-square test used for the differences variables between two or more categorical variables. Multivariate logistic regression analysis was used to establish a model to determine factors that are predictive of complicated appendicitis. The statistical significance was defined as  $P < 0.05$ .

## Results

Table 1 gives the comparison of sociodemographic and clinical characteristics of the appendicitis patients by gender. There were statistically significant differences between patients regarding age ( $P < 0.001$ ), BMI ( $P = 0.031$ ), cigarette smoking ( $P = 0.038$ ), sheesha smoking ( $P = 0.037$ ), family history of diabetes ( $P = 0.025$ ), hypertension ( $P = 0.019$ ), family history of gastrointestinal discomfort (GI) ( $P = 0.011$ ), and family history of appendicitis ( $P = 0.021$ ).

Table 2 shows the clinical characteristics symptoms' value among appendicitis by gender. Statistically significant differences were found between males and females for anxiety ( $P < 0.001$ ), red eye ( $P = 0.006$ ), dizziness ( $P = 0.021$ ), headache ( $P < 0.001$ ), muscular symptoms, weakness and cramps ( $P < 0.001$ ), bloating or swollen stomach ( $P < 0.001$ ), urinary tract infection (UTI) ( $P < 0.001$ ), chest pain ( $P < 0.001$ ), guarding ( $P < 0.001$ ), loss appetite ( $P = 0.004$ ), nausea ( $P < 0.001$ ) vomiting ( $P = 0.042$ ), anorexia ( $P = 0.009$ ), and constipation ( $P = 0.002$ ).

Table 3 presents the clinical sign and medical condition value among appendicitis by gender. There were statistically significant differences between males and females for pain ( $P < 0.001$ ), pain right belly ( $P = 0.027$ ), severe cramps ( $P = 0.007$ ), high temperature and fever ( $P < 0.001$ ), irritable bowel syndrome ( $P < 0.001$ ), RIF pain ( $P = 0.008$ ), rebound tenderness ( $P = 0.024$ ), positive bowel sounds ( $P = 0.029$ ), and pointing tenderness ( $P < 0.001$ ). Besides, Table 4 gives radiological diagnostic tests comparisons and their costs for appendicitis patients

Table 5 indicates multivariate stepwise logistic regression analysis of independent predictors for the presence of appendicitis and risk factors. Multivariate stepwise logistic regression analysis result showed nausea [3.46 (2.18–5.50)  $P < 0.001$ ]; C-reactive protein [2.95 (1.86–5.34)  $P < 0.001$ ]; dizziness [2.48 (1.18–5.20)  $P = 0.016$ ]; vomiting [2.37 (1.53–3.68)  $P < 0.001$ ]; muscular symptoms [1.98 (1.20–3.26)  $P = 0.007$ ]; irritable bowel syndrome [1.84 (1.55–218)  $P = 0.034$ ]; guarding [1.73 (1.44–3.36)  $P = 0.040$ ]; loss appetite [1.62 (1.19–2.60)  $P = 0.046$ ] were considered at higher risk as a predictors for appendicitis patients.

**Table 1: Comparison of sociodemographic and clinical characteristics of the patients by gender (n=643)**

Variables	Gender		P
	Males n=401	Females n=242	
Age groups (in years):			
20-29	152 (37.9)	47 (19.4)	0.001
30-39	108 (26.9)	58 (24.0)	
40-49	63 (15.7)	33 (13.6)	
50-59	43 (10.7)	54 (22.3)	
60 and above	35 (8.7)	50 (20.7)	
BMI (kg/m <sup>2</sup> )			
Normal (<25 kg/m <sup>2</sup> )	94 (23.4)	82 (33.9)	0.013
Overweight (29-30 kg/m <sup>2</sup> )	171 (42.6)	94 (38.8)	
Obese (>30 kg/m <sup>2</sup> )	136 (33.9)	66 (27.3)	
Physical activity			
Yes	111 (27.7)	53 (23.6)	0.248
No	290 (72.3)	185 (76.4)	
Smoking status			
Never	317 (79.1)	209 (86.4)	0.038
Current smoker	60 (15.0)	20 (8.3)	
Past smoker	24 (6.0)	13 (5.4)	
Sheesha smoking status			
Yes	69 (77.2)	27 (11.2)	0.037
No	332 (60.7)	215 (88.8)	
Family history of DM			
Yes	75 (18.7)	29 (12.0)	0.025
No	326 (81.3)	215 (88.0)	
Family history of hypertension			
Yes	94 (23.4)	38 (15.7)	0.019
No	307 (76.6)	204 (84.3)	
Family history of gastrointestinal discomfort (GI)			
Yes	71 (17.7)	25 (10.3)	0.011
No	330 (82.3)	217 (89.7)	
Family history of appendicitis			
Yes	72 (18.0)	27 (11.2)	0.021
No	329 (84.0)	215 (88.8)	

**Table 2: Clinical biochemistry baseline value and symptoms among appendicitis patients by gender (n=643)**

Variables	Males=401 n (%)	Females=242 n (%)	P
Anxiety	66 (16.5)	17 (7.0)	0.001
Red Eye	63 (15.7)	20 (8.3)	0.006
Dizziness	76 (19.0)	29 (12.0)	0.021
Headache	105 (26.2)	25 (10.3)	0.001
Muscular symptoms, weakness	87 (21.7)	27 (11.2)	0.001
Bloating/swollen stomach	75 (18.7)	18 (7.4)	0.001
Urinary tract infections -UTI	68 (17.0)	16 (6.6)	0.001
Chest pain	53 (13.2)	12 (5.0)	0.001
Guarding	77 (19.2)	18 (7.4)	0.001
Loss appetite	96 (23.9)	35 (14.5)	0.004
Nausea	96 (23.9)	31 (12.8)	0.001
Vomiting	108 (26.9)	48 (19.8)	0.042
Anorexia	80 (20.0)	29 (12.0)	0.009
Constipation	90 (22.4)	30 (12.4)	0.002
<b>Biochemistry</b>			
Parameters	Mean±SD	Mean±SD	P
C-reactive protein - CRP (mg/L)	37.4±13.9	34.3±16.4	0.002
White Blood Count (/mL)	13840.1±5,346.5	12,528.5±4,864.2	0.005
Systolic blood pressure (mmHg)	128.5±15.1	125.1±12.4	0.001
Diastolic blood pressure (mmHg)	80.4±9.3	78.2±9.1	0.002

**Table 3: Clinical biochemistry baseline value among appendicitis patients by gender (n=643)**

Variables	Males n=401 n (%)	Females n=242 n (%)	P
Pain	134 (33.4)	45 (18.6)	0.001
Pain right belly	46 (11.5)	15 (6.2)	<0.027
Pain left belly	45 (11.2)	17 (7.0)	<0.081
Severe cramps	116 (28.9)	47 (19.4)	0.007
High temperature-fever	67 (16.7)	19 (7.9)	0.001
Painful peeing	50 (12.5)	28 (11.6)	0.735
Irritable bowel syndrome	69 (17.2)	20 (8.3)	0.001
RIF Pain	64 (16.0)	21 (8.7)	<0.008
Rigidity	72 (18.0)	36 (14.9)	0.312
Rebound tenderness	53 (13.2)	18 (7.4)	0.024
Positive bowel sound	82 (20.4)	33 (13.6)	0.029
Obturator Sign	55 (13.7)	28 (11.6)	0.432
Psoas Sign	46 (11.5)	26 (10.7)	0.777
Rovsing's Sign	88 (17.0)	32 (13.2)	0.206
Percussion Tenderness	42 (10.5)	15 (6.2)	<0.065
Pointing Tenderness	58 (14.5)	14 (5.8)	0.001

**Table 4: Radiological diagnostic test and their costs for appendicitis patients**

Patient Group	Appendicitis	
	Number	%
Compliant population		
Ultrasound	185	28.8
Computed tomography	298	46.3
Ultrasound and computed tomography	160	26.9
Radiological Test cost	Price TL	Price \$-US Dollar
National Health Insurance	5,500 TL	\$1,000
Private Insurance	8000-12000 TL	\$1,500-\$2,000
Non-Insurance	8,000 TL	\$1,500
Physician exam cost	800 TL	\$150

TL=Turkish Lira and \$1=0.5500 TL.

**Table 5: Multivariate stepwise logistic regression analysis of independent predictors for the appendicitis**

Variables	Odds ratio (95%CI)	P
Nausea	3.46 (2.18-5.50)	<0.001
C-reactive protein - CRP (mg/L)	2.95 (1.86-5.34)	<0.001
Vomiting	2.37 (1.53-3.68)	<0.001
Muscular symptoms	1.98 (1.20-3.26)	0.007
Dizziness	2.48 (1.18-5.20)	0.016
Irritable bowel syndrome	1.84 (1.55-218)	0.034
Guarding	1.73 (1.44-3.36)	0.040
Loss Appetite	1.62 (1.19-2.60)	0.046

## Discussion

The clinical diagnosis of acute appendicitis in the early phases of the disease is difficult as it may mimic other conditions. The newer techniques of US and CT have shown great promise in evaluation of patients with suspected acute appendicitis.

On patients suspected to have acute appendicitis admitted to the primary care institution, the US and CT should be used for

diagnosis. Diagnosed and suspected patients should be directed to a surgical center.

However, advantages and limitations exist in both US and CT for evaluating patients with suspected acute appendicitis. In the current study, the US was performed on 185 (28.8%) patients, CT conducted on 298 (46.3%) patients, and 160 (26.9%) performed on both US and CT for diagnosing appendicitis. The outcome results are comparable and consistent with the previously reported studies.<sup>[12-16]</sup> Generally, CT is widely accepted and the preferred modality for evaluation of suspected appendicitis because of its great diagnostic performance,<sup>[19,20]</sup> speed and good interobserver agreement for interpretation regardless of experience. We were able to identify essential risk factors and predictors based on these images that can be used to assign a high probability of appendicitis in the US and CT.

Acute appendicitis is the most common abdominal surgical emergency that can affect individuals from all age groups. The prevalence of appendicitis in the current study occurred higher among young age groups 20–39 years old 48.0% among males and 43.4% among females and this confirmative with previous report study in United States age groups 18–39 years old by 55.4%.<sup>[21]</sup> The present study revealed that the prevalence of appendicitis is higher among males (62.6%) compared to the females (37.6%), this is consistent with the previously reported appendicitis prevalence by gender in France<sup>[22]</sup> (males 57.8% vs females 42.2%). Moreover, the increased risk of male versus female and age <50 versus age > is in line with the recent literature<sup>[23]</sup> and confirming our study.

An accurate diagnosis of acute appendicitis can be established with great confidence in the majority of patients, based on history, and physical examination. The present study revealed that pain, anorexia, vomiting, nausea, temperature >37.3°C, rebound tenderness, percussion tenderness, white cell count >10 × 10<sup>9</sup>/L, loss appetite, constipation, and severe cramps were common significant risk factors among patients.<sup>[3,8,12-16,24]</sup>

A family history of acute appendicitis is an important factor determining the likelihood of appendicitis and must be considered during the medical visit. Clinicians attempting to confirm their diagnostic accuracy when patients present with acute abdominal pain should inquire about family history of appendicitis. Gauderer *et al.*<sup>[17]</sup> suggested that children who have appendicitis are twice more likely to have a positive family history than are those with right lower quadrant pain. The complex segregation analysis supported a polygenic or multifactorial model with a total heritability of 56%<sup>[25]</sup> among appendicitis patients.

## Limitations and strength of the study

Our study has several limitations. Firstly, the sample might be partially biased due to the consecutive series of patients with the prospective cohort study. Secondly, we did not have data

on family history in our study population. Hence, our results relied solely on the patients' knowledge of their family history response. Thirdly, the gender proportion of males and females were not balanced. Finally, no pathological results were available for some patients.

## Conclusion

In conclusion, CT offers the best cost-effectiveness in the prepaid system and public health system. The current study suggested that nausea, C-reactive protein, dizziness, vomiting, muscular symptoms, irritable bowel syndrome, guarding, and loss appetite were considered at higher risk as a predictor for appendicitis patients.

## Acknowledgments

This work was supported by the Istanbul Medipol University, International School of Medicine. The authors would like to thank the Istanbul Medipol University for their support and ethical approval (Research Protocol and IRB# 10840098-604.01.01-E.45424).

## Financial support and sponsorship

Nil.

## Conflicts of interest

There are no conflicts of interest.

## References

- Öztürk A, Yananlı Z, Atalay T, Akıncı ÖF. The comparison of the effectiveness of tomography and Alvarado scoring system in patients who underwent surgery with the diagnosis of appendicitis. *Ulus Cerrahi Derg* 2015;32:111-4.
- Alireza P, Fatemeh S, Naghme R. Diagnostic accuracy of abdominal ultrasonography in pediatric acute appendicitis. *Bull Emerg Trauma* 2019; 7:278-83.
- Johansson EP, Rydh A, Riklund KA. Ultrasound, computed tomography, and laboratory findings in the diagnosis of appendicitis. *Acta Radiol* 2007;48:267-73.
- Leite PN, Pereira JM, Cunha R, Pinto P, Sirlin C. Computed tomography evaluation of appendicitis and its complications: Imaging techniques and key diagnostic findings. *Am J Roentgenol* 2005;185:406-17.
- Parks NA, Schroepel TJ. Update on imaging for acute appendicitis. *Surg Clin North Am* 2011;91:141-54.
- SCOAP Collaborative, Cuschieri J, Florence M, Flum DR, Jurkovich GJ, Lin P, *et al*. Negative appendectomy and imaging accuracy in the Washington state surgical care and outcomes assessment program. *Ann Surg* 2008;248:557-63.
- Raja AS, Wright C, Sodickson AD, Zane RD, Schiff GD, Hanson R, *et al*. Negative appendectomy rate in the era of CT: An 18-year perspective. *Radiology* 2010;256:460-5.
- Gamanagatti S, Vashisht S, Kapoor A, Chumber S, Bal S. Comparison of graded compression ultrasonography and unenhanced spiral computed tomography in the diagnosis of acute appendicitis. *Singapore Med J* 2007;48:80-7.
- Long SS, Long C, Lai H, Macura KJ. Imaging strategies for right lower quadrant pain in pregnancy. *AJR* 2011;196:4-12.
- Toorenvliet BR, Wiersma F, Bakker RF, Merkus JW, Breslau PJ, Hamming JF. Routine ultrasound and limited computed tomography for the diagnosis of acute appendicitis. *World J Surg* 2010;34:2278-85.
- Krishnamoorthi R, Ramarajan N, Wang NE, Newman B, Rubesova E, Mueller CM, *et al*. Effectiveness of a staged US and CT protocol for the diagnosis of pediatric appendicitis: Reducing radiation exposure in the age of ALARA. *Radiology* 2011;259:231-9.
- Kim MS, Kwon H-J, Kang KA, Do IG, Park HJ, Kim EY, *et al*. Diagnostic performance and useful findings of ultrasound re-evaluation for patients with equivocal CT features of acute appendicitis. *Br J Radiol* 2018;91:20170529.
- Sim JY, Kim HJ, Jang SK, Yeon JW, Jeon BG, Ha YR, *et al*. Value of additional ultrasound examination in patients with equivocal computed tomography findings of acute appendicitis: Comparison with computed tomography reassessment. *J Med Ultrasound* 2019;27:75-80.
- Franke C, Bohner H, Yang Q, Ohmann C, Roher HD. Ultrasonography for diagnosis of acute appendicitis: Results of a prospective multicenter trial. *World J Surg* 1999;23:141-6.
- Pickuth D, Heywang-Kobrunner SH, Spielmann RP. Suspected acute appendicitis: Is ultrasonography or computed tomography the preferred imaging technique? *Eur J Surg* 2000;166:315-9.
- Wise SW, Labuski MR, Kasales CJ, Blebea JS, Meilstrup JW, Holley GP, *et al*. Comparative assessment of CT and sonographic techniques for appendiceal imaging. *AJR Am J Roentgenol* 2001;176:933-41.
- Gauderer MWL, Crane MM, Green JA, Decou JM, Abrams RS. Acute appendicitis in children: The importance of family history. *J Ped Surg* 2001;36:1214-7.
- Bennington-Castro J, Jasmer R. Symptoms of Appendicitis: Nausea, Fever, Abdominal Pain, and More Stomach Pain. Available from: <https://www.everydayhealth.com/appendicitis/guide/symptoms/>. [Last accessed on 2020 Jan 18].
- Bener A, Brebner J, Norman JN, El-Ghazawi I, Al-Suwaidi MHMO, Abu-Azab I. Remote general practice: Diagnosis of appendicitis. *Can J Rural Med* 2002;7:26-9.
- Van den Worm L, Georgiou E, De Klerk M. C-reactive protein as a predictor of severity of appendicitis. *S Afr J Surg* 2017;55:14-7.
- Drake FT, Mottey NE, Farrokhi ET, Florence MG, Johnson MG, Mock C, *et al*. Time to appendectomy and risk of perforation in acute appendicitis. *JAMA Surg* 2014;149:837-44.
- Gignoux B, Blanchet MC, Lanz T, Vulliez A, Saffarini M, Bothorel H, *et al*. Should ambulatory appendectomy become the standard treatment for acute appendicitis? *World J Emerg Surg* 2018;13:28.
- Lally KP, Cox CS, Andrassy RJ. Appendix. In: Townsend MC Jr, Beauchamp RD, Evers BM, Mattox KL, editors *Text Book of Surgery*. 17<sup>th</sup> ed. London: WB Saunders; 2004. p. 1381-99.
- Ergul E, Ucar AE, Ozgun YM, Korukluoglu B, Kusdemir A. Family history of acute appendicitis. *J Pak Med Assoc* 2008;58:635-7.
- Simó Alari F, Gutierrez I, Giménez Pérez J. Familial history aggregation on acute appendicitis. *BMJ Case Rep* 2017;2017.