

Research Article

Nihal Karakaş*, Mehmet Evren Okur, Nurşah Öztunç, Derya Çiçek Polat and Ayşe Esra Karadağ



Laurocerasus officinalis Roem. fruit extract induces cell death through caspase mediated apoptosis in gastric cancer cell lines

Laurocerasus officinalis Roem. meyve ekstralarının mide kanseri hücrelerinde kaspaz aracılı apoptoz yoluyla sitotoksik etkilerinin incelenmesi

<https://doi.org/10.1515/tjb-2020-0378>

Received February 23, 2020; accepted October 22, 2020; published online January 8, 2021

Abstract

Objectives: *Laurocerasus officinalis* Roem. fruits are traditionally used for several health problems. Although there are some studies about its antiproliferative effects on different cancer cells, no study was reported about its potential therapeutic efficacy against gastric cancers which is the most malignant disease in the digestive system with high morbidity and mortality.

Methods: This study was aimed to evaluate *L. officinalis* fruit extract phytochemical contents as well as to compare

anticancer effects on gastric cancer cells. The antioxidant activities were determined by ABTS and DPPH assays. Anti-cancer effects were measured by cell viability assays, then apoptotic proteins were analyzed by western blotting and flow cytometry.

Results: *Laurocerasus officinalis* fruit methanol extract showed moderate antioxidant activity by ABTS* and DPPH* assays. Significant cytotoxic activities and caspase mediated apoptosis were detected in the extract treated MKN-45 and AGS gastric cancer cells respectively while sparing healthy cells.

Conclusion: Our results showed that the *L. officinalis* Roem. extract has significant anticancer efficacy on gastric cancer cell lines; therefore, it can be further studied to determine its potential therapeutic components.

Keywords: antioxidant activity; apoptosis; gastric cancer; *Laurocerasus officinalis*; Rosaceae.

Öz

Giriş: *Laurocerasus officinalis* Roem. meyveleri halk arasında çeşitli sağlık problemlerine karşı kullanılmaktadır. Farklı kanser hücreleri üzerindeki antiproliferatif etkileri ile ilgili bazı çalışmalar olmakla birlikte, sindirim sistemindeki ciddi bir rahatsızlık olan, yüksek morbidite ve mortaliteye sahip mide kanserlerine karşı herhangi bir çalışma bildirilmemiştir.

Yöntem: Bu çalışma *L. officinalis* meyve ekstralarının fitokimyasal analizi ile beraber karşılaştırmalı olarak mide kanseri hücrelerine karşı sitotoksik etkilerinin incelenmesini amaçlamıştır. Antioksidan aktivite ABTS ve DPPH yöntemleriyle araştırılmıştır. Antikanser etkinliği ise hücre canlılığı yöntemi ile araştırılmış ve sonrasında apoptotik

*Corresponding author: Nihal Karakaş, Department of Medical Biology, School of Medicine, Istanbul Medipol University, Beykoz, Istanbul, Turkey; and Research Institute for Health Sciences and Technologies (SABITA), Istanbul Medipol University, Istanbul, Turkey, Phone: +90 216 681 51 00, E-mail: nkarakas@medipol.edu.tr, <https://orcid.org/0000-0002-9096-1512>

Mehmet Evren Okur, Department of Pharmacology, Faculty of Pharmacy, University of Health Sciences, Istanbul, Turkey, E-mail: evrenokurecz@gmail.com

Nurşah Öztunç, Research Institute for Health Sciences and Technologies (SABITA), Istanbul Medipol University, Istanbul, Turkey

Derya Çiçek Polat, Department of Pharmaceutical Botany, Faculty of Pharmacy, Ankara University, Ankara, Turkey, E-mail: polatd@ankara.edu.tr

Ayşe Esra Karadağ, Department of Pharmacognosy, School of Pharmacy, Istanbul Medipol University, Istanbul, Turkey; and Graduate School of Health Sciences, Anadolu University, Eskişehir, Turkey, E-mail: ayseesraguler@gmail.com

proteinleri western blot ve flow sitometre yöntemleri ile ölçülmüştür.

Sonuçlar: *L. officinalis* meyve metanol ekstresi, ABTS ve DPPH testlerine göre orta düzeyde antioksidan aktivite göstermiştir. Sağlıklı hücrelerde sitotoksik etki gözlemlenmezken, ekstre uygulanmış MKN-45 ve AGS mide kanseri hücrelerinde önemli sitotoksik aktiviteler ve kaspaz aracılı apoptoz saptanmıştır.

Tartışma: Sonuçlar, *L. officinalis* Roem ekstresinin mide kanseri hücre hatları üzerinde önemli antikanser etkinliğine sahip olduğunu göstermiştir; bu nedenle, potansiyel terapötik bileşenlerini belirlemek için ileri çalışmalar gerçekleştirilebilecektir.

Anahtar kelimeler: *Laurocerasus officinalis*; Rosaceae; mide kanseri; apoptoz; antioksidan aktivite.

Introduction

Laurocerasus officinalis Roem. (synonym: *Prunus laurocerasus* L.) is a food plant of the Rosaceae family. It grows naturally in Turkey and is known as “laz kirazi, karayemiş or taflan”. Cherry laurel is native of Central and Western Asia, Anatolia, and Southern Europe and cultivates in temperate regions and is mostly used as ornamental plant [1]. Its fruits are usually consumed as jam, marmalade, fruit juice, tea, and in canned or pickled styles [2]. Fruits and seeds are used in the treatment of kidney stones, stomach ulcers, bronchitis, strengthening the bones, the acid-base balance of blood (seeds), eczemas and hemorrhoids, and as a diuretic, antispasmodic, antitussive (fruits) as a folk medicine in Turkey [3]. The major bioactive components are 2-O-β-d-glucopyranosyl-2-hydroxyphenyl-acetic acid, (+)-catechin and kaempferol-3-O-β-d-xylopyranosyl-(1→2)-O-β-d-glucopyranoside have been noticed in the cherry laurel leaves extract [4]. *L. officinalis* fruit stems nutritional and pharmaceutical value from its vanillic, caffeic, chlorogenic, and benzoic acid with fructose, glucose [5], mannitol, ascorbic acid, antocyanins [6] and tannin content [7].

Although many drugs have been developed for the treatment of cancers, there are concerns about the therapeutic effects and safety of these drugs. The major problem of chemotherapeutic drugs used as a standard treatment in various types of cancers is the toxicity [8]. However, products from plants have been proven to be effective and safe in the treatment of cancers. Therefore, cancer drug discoveries are also directed to plant derived products obtained from natural plants [9]. These products act as anti-cancer agents by interfering with the initiation, development, and progression of cancer through the

modulation of various mechanisms including cellular proliferation, differentiation, apoptosis, angiogenesis, and metastasis [10]. Studies have greatly agreed on the protective effect of fruits and vegetables in reducing the risk of stomach cancer, because fruits and vegetables contain the antioxidant substances [11–13].

Laurocerasus officinalis is used for stomach ulcer treatment as a folk medicine in Turkey and therefore its contents make it a potential therapeutic candidate for different gastrointestinal problems.

Gastric cancer is a malignant disease and ranks fifth for cancer incidence and second for cancer deaths [14]. The underlying mechanisms of gastric cancer are mostly linked to *Helicobacter Pylori* (*H. Pylori*) infection and abundance of the *H. Pylori* makes its mortality pattern variative for different ethnics. Eradication of *H. Pylori* provides a good prognosis of the disease [15–17].

Unfortunately, the infection is hard to clear by the host and results in a chronic inflammatory state with continued oxidative stress within the tissue. Reactive oxygen/nitrogen species are released by the effected immune and epithelial cells; then damage the surrounding tissue and lead to gastric carcinogenesis [18]. Therefore, *H. Pylori* behavior and host response determine the progress of the disease, and alternative treatment options are needed to overcome chemotherapy resistance developed by the bacterium [19].

There have been many anti-cancer studied on *L. officinalis* and showed a selective cytotoxic effect in lung, colon, prostate, liver, and cervical cancer cell lines [20–22]. But no studies have been found with anti-cancer activities against gastric cancer *L. officinalis*. In this study, we focused on its potential anti-cancer effects against gastric cancer since it can be eaten and in direct contact with the stomach. Besides cell viability assays, we showed apoptotic cell death by western blotting and Annexin V/PI stainings upon treatment with the extract.

In the present study, *L. officinalis* fruits total phenolic content and flavonoid contents were determined, afterward; their *in vitro* antioxidant and anti-cancer activities against gastric cancer showed that it can be further studied and characterized for therapeutic benefits.

Materials and methods

Materials

The standard chemicals were purchased from Sigma Chemical Co. (St. Louis, USA) and the HPLC-grade solvents were purchased from Merck. Methanol was purchased from Sigma-Aldrich, Germany. All other reagents and solvents used were of analytical grade.

Preparation of samples

Laurocerasus officinalis fruits were collected from Ankara University, Ankara, Turkey (Date: 18.07.2017), and identified by Derya Çiçek Polat. *L. officinalis* fruit, which is dark purple, were pureed and extracted with methanol on a magnetic stirrer (Heidolph MR3001, Germany) (250 g sample, 400 mL × 3 days) followed by filtration [1]. The extract was distilled using an evaporator (Heidolph WB2000, Germany).

Total phenolic and flavonoid contents

Folin Ciocalteu technique was used to detect total phenolics of the fruit extract. The mixture was prepared with fruit extract (5 mL), Folin-Ciocalteu's reagent (0.25 mL), and Na₂CO₃ (0.2 mL) and kept for 15 min at 45 °C. The absorbance reading of samples was performed at 765 nm. A calibration curve (R²=0.981) was used for calculating the total phenolic content (TPC) of the extract and the outcome was showed as mg gallic acid equivalent (gae)/100 g extract [1, 23, 24]. The basis of this method is based on the redox reaction of phenolic compounds in the basic medium, reducing the Folin-Ciocalteu reagent and converting the phenolic compounds into oxidized form. The color intensity of the resulting complex is directly proportional to the concentration of phenolic substances [25].

The aluminum chloride colorimetric method was performed to detect the total flavonoid content of *L. officinalis* fruit extract. 50 µL of the extract was mixed with methanol (up to 1 mL) and added 4 mL water, then 5% NaNO₂ solution; 10% AlCl₃ solution was added. Afterward, NaOH (1 mol/L) was added and water was used to adjust to 10 mL (total volume). After waiting, the absorbance of the mix was read at 510 nm. The content of flavonoid was detected by the calibration curve (R²=0.9978) [26] and the outcome was displayed as mg quercetin equivalent (qe)/100 g extract. All experiments were done in triplicate. The basis of this method is that Aluminum chloride forms acid stable complexes with the C-4 keto groups and either the C-3 or C-5 hydroxyl group of flavonoids. Moreover, aluminum chloride is also complex with ortho-dihydroxyl groups of A or B-rings of flavonoids. The results are usually given as equivalent to standard flavonoids quercetin [27].

DPPH scavenging assay

To determine the *L. officinalis* fruit extract antioxidant ability, the DPPH (2,2-diphenyl-1-picrylhydrazyl) radicals were utilized according to the spectrophotometric protocol [1, 28]. The absorbance was read at 517 nm using a spectrophotometer. The radical scavenging activity was calculated according to the following equation:

$$\text{DPPH-RSA \%} = \left(\frac{[\text{Absorbance control} - \text{Absorbance test sample}]}{[\text{Absorbance control}]} \right) \times 100$$

All experiments were done in triplicate. Ascorbic acid was served as a positive control. IC₅₀ rates were detected from a calibration curve [23].

ABTS scavenging assay

2,2'-azino-bis-3-ethylbenzthiazoline-6-sulfonic acid (ABTS) radical scavenging activity of *L. officinalis* was determined according to Re et al. [29]. Stock ABTS solution was composed by reacting aqueous of

ABTS with potassium persulfate solution. The mixed solution was incubated for 12–16 h in the dark at room temperature. The absorbance of reaction mixtures was measured at 734 nm. Three independent experiments were performed. An analog of vitamin E, Trolox was used as the positive control [1, 26]. The results were compared with Trolox and expressed as IC₅₀ as follows:

$$\% \text{ ABTS inhibition} = \left(\frac{[\text{Absorbance control} - \text{Absorbance test sample}]}{[\text{Absorbance control}]} \right) \times 100$$

LC-MS analysis

For Liquid Chromatography Mass Spectroscopy (LC-MS) analysis the extract was dissolved in methanol and filtered. The extract was analyzed using LC-MS (1200LC, Agilent). Ionization was achieved in negative mode with ESI. For the chromatographic separation, LC-MS was run on 4.6 × 250 mm, i.d. 5 µm particle size, octadecyl silica gel analytical C18 column, and its temperature was maintained at 40 °C. The elution gradient consisted of mobile phases were A: Acetonitrile:Water (10:90, v/v) and B: Acetonitrile:Water (90:10 v/v). The gradient established in the time frame 0–40 min, B% 15–100. The solvent flow rate is maintained at 0.7 mL/min. The injection volume is 20 µL. The range of 100–1000 amu was scanned and recorded for MS analysis [30].

Cell culture

AGS-human gastric adenocarcinoma (ATCC, #CRL-1739) cell line and Human Primary Dermal Fibroblasts (HDFa) (ATCC, # PCS-201-012) were purchased from ATCC (U.S.). The human gastric adenocarcinoma cell line, MKN-45 (DSMZ, #ACC409) was obtained from DSMZ (Germany). AGS and MKN-45 cells were grown in RPMI and DMEM (Gibco) medium respectively in the presence of 10% fetal bovine serum (Gibco) with the addition of 1% antibiotics (penicillin/streptomycin) in 5% CO₂ incubator (at 37 °C). The subculture of cells was performed every 4–5 days as the cells reach the confluency. The cells were harvested from the flask with Trypsin/EDTA 0.25% (Gibco) and for cell viability assays, 5 × 10³ cells/well were seeded into 96 black well plates (Corning).

Cell viability assays

Extracts were dissolved in growth medium and filtered using 0.20 µm PES filters (Sarstedt) to prepare stock solutions, and for serial dilutions as a final concentration to normalize measurements. After seeding into 96 well plates, for 24 h, cells were incubated in 5% CO₂ (37 °C) incubator. Then, 1–10 mg/mL of *L. officinalis* extract was added as triplicates. After 48 h of treatment, Cell Titer Glo reagent (Promega) added into each well according to the manufacturer's guide, and viable cells were determined by reading luminescence signal under SpectraMax i3x Multi-Mode Detection Platform.

Western blotting

Cells were seeded at a density of 2 × 10⁵ cells/well into six well plates for western blot sample collection. Then incubated at 37 °C in 5% CO₂ for 24 h. The culture medium was discarded, and cells were treated with 0 mg/mL (control) or 5 mg/mL of *L. officinalis* extract (Control wells were treated with an equal amount of extract solvent; dH₂O). After 48 h of

treatment, protein lysates were obtained using Ripa lysis buffer (Thermo Fischer Scientific; #89900) from each well. Protein samples were equally loaded (25 µg/well) and run on SDS-PAGE. Then, Bio-Rad semi-dry western blotting protocol was applied. Then the membrane was incubated with blocking buffer (5% BSA or 5% skim milk accordingly) for 1 h at room temperature. Afterward, the membrane was probed with antibodies against PARP, caspase-3 and β-actin (anti-PARP (CST; #9542), anti-cleaved caspase 3 (CST; #9161), (CST #4970), anti-rabbit (CST; #7074) and HRP conjugated anti-mouse (GenDEPOT; #W3903) antibodies were used). The first antibody incubation was performed at 4 °C for overnight and after washing with TBST three times, membranes were then probed with HRP conjugated secondary antibodies for 2 h at room temperature. The membranes were washed and incubated with a 1:1 ratio of Clarity Western ECL Substrate (Bio-Rad), then analyzed for protein bands by ChemiDoc-MP (Bio-Rad).

Annexin V/Propidium Iodide apoptosis stainings

After seeding into 100 × 20 mm culture dishes, cells were incubated at 37 °C in 5% CO₂ for 24 h. Then the culture medium was discarded, and cells were treated with 0 mg/mL (control) and 5 mg/mL, of *L. officinalis* extract. To detect early apoptotic cells, an earlier time point was determined than the viability assays. Therefore, after 36 h of treatment with *L. officinalis* extract, Annexin V-FITC/Propidium Iodide (PI) early apoptosis double staining protocol was applied according to manufacturer's instructions (CST #6592 Annexin V-FITC Early Apoptosis Kit). Then, early apoptotic, late apoptotic, necrotic, and live cell percentages were determined through flow cytometry analysis.

Statistical analysis

All statistical comparisons of antioxidant activity studies were performed by one-way ANOVA followed by Dunnett's tests. $p < 0.05$ was considered statistically significant. For *in vitro* cell based assays, statistical comparisons were performed by unpaired Student's *t*-test assuming equal variance. Differences were considered as statistically significant at $0.003 < p^* \leq 0.005$ and $0.0005 < p^{**} \leq 0.003$; $p^{***} \leq 0.0003$ and $0.01 < p^{\#} \leq 0.05$. Data were expressed as means ± S.E.M.

Results

Phenolic and flavonoid contents

Total phenolics were calculated by using the Folin-Ciocalteu method. Total flavonoids in the fruit extract were measured using the aluminum chloride colorimetric method. *L. officinalis* fruit total phenolic and flavonoid content were detected, and the results which are similar to our previous study [1] are given in Table 1.

In vitro antioxidant assays

The free radical scavenging activity of the extract was determined using ABTS and DPPH experiments and the

Table 1: Total flavonoid and phenolic content of *L. officinalis* fruit extract.

	Total flavonoid content mg QE/100 g extract	Total phenolic content mg GAE/100 g extract
<i>L. officinalis</i> fruit extract	502.10 ± 6.85 mg qe/100 g	461.31 ± 4.98 mg gae/100 g

outcomes are assembled in Table 2, as previously reported [1]. The antioxidant activity of *L. officinalis* was investigated by similar and different methods. When the results were compared, similar results were observed in similar studies. It is known that the collection of the plant, its drying, the area it was grown, and the extraction method even caused a change in the antioxidant capacity. Therefore, some minor differences can be observed between previous studies [6, 31–34].

Qualitative chromatographic analysis by LC-MS

The flavonoid compositions of the methanol extract were detected by LC/MS. Standards were detected according to mass analyses and qualitative analyses using reversed-phase chromatography. The flavonoid components of *L. officinalis* fruit were characterized as luteolin, quercetin, and apigenin (Figure 1, Figure 2) (Table 3).

Cell viability assays

Treatment with *L. officinalis* fruit extract led to significantly decreased cell viability in MKN45 and AGS human gastric cancer cell lines

Cytotoxic effects of the *L. officinalis* on MKN45 (human gastric cancer cell line), AGS (gastric adenocarcinoma cell line), and primary human fibroblast cells by measuring their metabolically active state. According to our findings, cell viability significantly decreased when treated with

Table 2: ABTS and DPPH scavenging activities of *L. officinalis* extract.

	<i>L. officinalis</i> extract		References
	IC50 ± SD, mg/mL		
ABTS	2.44 ± 0.21		2.986 (trolox)
DPPH	2.95 ± 0.09		3.767 (ascorbic acid)




Figure 1: *L. officinalis* fruit methanol extract LC-MS chromatogram.




Figure 2: Standard chromatogram of LC-MS. Standards:1, gallic acid (R.T. 4.96); 2, Luteolin-7-o-glycoside (R.T. 7.62); 3, Coumaric acid (R.T. 9.21); 4, Rosmarinic acid (R.T. 10.65); 5, Myrcetin (R.T. 11.43); 6, Quercetin (R.T. 15.25); 7, Apigenine (R.T. 18.47); 8 Luteolin (R.T. 19.15) (R.T.=Retention time).

Table 3: *L. officinalis* methanol extract flavonoid compounds.

Compounds	RT	Base peak (m/z)
Quercetine	15.418	314.99
Apigenine	18.502	269.12
Luteolin	19.213	285.06

5–10 mg/mL extract in both MKN-45 and AGS cancer cell lines. Treatment of gastric cancer cell lines with 5 mg/mL *L. officinalis* extract showed 65.6% ($p=0.0031$) and 25.2% ($p=0.0038$) significantly decreased cell viability in AGS and MKN-45 cells, respectively. Besides that, 10 mg/mL extract of *L. officinalis* led to highly decreased viability of cells with 1.1% in MKN45 and 27.4% in AGS cell lines ($p \leq 0.0003$) Conversely, treatment of primary human fibroblast cells with the same concentrations (5–10 mg/mL) of *L. officinalis* extracts showed statistically non-significant cytotoxicity (Figure 3A). Accordingly, *L. officinalis* extract showed 3.770 mg/mL; 5.606 mg/mL and 29.42 mg/mL IC_{50}

values for MKN-45; AGS and primary human fibroblast cells respectively (Figure 3B).

Detection of apoptosis

L. officinalis extract induced apoptosis in both AGS and MKN-45 cell lines through caspase-3 and PARP cleavages

Apoptosis was analyzed in 5 mg/mL extract treated MKN-45 and AGS gastric cancer cell lines. Compared to control treatments (0 mg/mL), cleaved PARP and cleaved caspase-3 presence were detected in *L. officinalis* extract treated AGS and MKN-45 gastric cancer cell lines (Figure 4A). According to Image J analysis of AGS cell line, cleaved PARP intensity increased approximately four folds ($p \leq 0.0001$), and cleaved caspase three intensity increased approximately eight folds ($p=0.0025$) (Figure 4B). In *L. officinalis* extract treated MKN-45 cell




Figure 3: Cytotoxic effects of *L. officinalis* extracts on MKN45 and AGS gastric cancer cell lines and primary human fibroblast cells. Cell viability of a) AGS and MKN-45 cell line decreased in a concentration dependent manner upon 48 h treatment with 1–10 mg/mL *L. officinalis* extracts. Decrease in viability of fibroblast cells were found statistically not significant. Cell viability assays were performed as three independent experiments. Data are expressed as \pm SEM. Differences were considered as statistically significant at $0.003 < p \leq 0.005$ and $0.0005 < p^{**} \leq 0.003$; and $p^{***} \leq 0.0003$. b) Normalized dose vs. response graphs and IC₅₀ values of AGS; MKN45 cell lines and primary human fibroblast cells. Accordingly, *L. officinalis* extract showed IC₅₀ values of 3.770 mg/mL in MKN45 cell line; 5.606 mg/mL in AGS cell line; and 29.42 mg/mL in primary human fibroblast cells.

line; cleaved PARP intensity increased approximately one and a half folds ($p=0.0029$) and cleaved caspase-3 intensity increased approximately five folds ($p=0.0010$) (Figure 4C).

These data demonstrated the cleavage of apoptotic proteins and biochemical verification of cell death induced by *L. officinalis* extract in both AGS and MKN-45 gastric cancer cell lines. Additionally, caspase cleavage showed that *L. officinalis* extract induced cell death occurs in a caspase dependent manner.

Apoptotic cell populations were determined in *L. officinalis* fruit extract treated AGS cell line

We analyzed the apoptotic phase of cell populations in *L. officinalis* extract treated AGS gastric adenocarcinoma cell line. Different from cell viability assays, AGS cells were either treated with distilled water and 5 mg/mL of *L. officinalis* just for 36 h to track early apoptotic cells as well as late apoptotic cells using Annexin V-FITC Early Apoptosis Detection Kit (CST, U.S.). According to staining, *L. officinalis* extract treated AGS cells showed a shift from alive to death cell state, which is significantly determined as apoptotic cells either

from early or late stages (Figure 5A, B). The quantitative analysis was performed and when compared to control cells, viable cells decreased from 89.65 to 66.22%, and early apoptotic cells increased from 1.55 to 19.77%, while late apoptotic cells increased from 4.78 to 11.78%. Necrotic cell increase in population was not found statistically significant (from 4.02 to 2.23%) (Figure 5C).

Discussion

In this study, we evaluated the potential use of *L. officinalis* fruit methanol extract against gastric cancer cell lines as it is one of the most malignant cancers studied and traditionally used in Turkey for the treatment of different types of human illness, especially for gastrointestinal problems [6, 33, 35, 36]. Firstly, *L. officinalis* fruits TPC and TFC were determined and they have been investigated with respect to their *in vitro* antioxidant and anti-cancer activities against gastric cancer.

In our study, we found the total phenolic compounds and total flavonoids which are responsible for antioxidant activity were also rich in fruits.

The antioxidant activities of fruits were measured using the ABTS and DPPH methods which are most




Figure 4: Western blot analysis of cell death in *L. Officinalis* fruit extract treated AGS and MKN45 gastric cancer cell lines. a) AGS and MKN45 gastric cancer cell lines were treated with distilled water (0 mg/mL) (left band) and 5 mg/mL of *L. Officinalis* (right band) for 48 h to detect apoptosis. In comparison with control bands (0 mg/mL), cleaved PARP and cleaved caspase three presence were detected in 5 mg/mL treatments. According to Image J analysis of b) AGS and c) MKN-45 gastric adenocarcinoma cell lines; cleaved PARP and cleaved caspase-3 bands were detected as compared to control treatments. Data are expressed as \pm SEM. Differences were considered as statistically significant at $0.0005 < p^{**} \leq 0.003$; and $p^{***} \leq 0.0003$. Experiments were performed as triplicates and band densities were normalized to actin controls using image J analysis.




Figure 5: Flow cytometry analysis of apoptosis in 5 mg/mL *L. officinalis* treated AGS gastric adenocarcinoma cell line. AGS cells were either treated with a) distilled water (left) or b) 5 mg/mL of *L. officinalis* (right) for 36 h to detect early apoptotic cells using Annexin V-FITC Early Apoptosis Detection Kit (CST, U.S.). Panels P2-Q1; P2-Q2; and P2-Q3 indicate necrotic cells; late apoptotic cells, and early apoptotic cells respectively. Each staining was performed as triplicates. Data are expressed as \pm SEM and differences were considered as statistically significant at $0.01 < p^{\#} \leq 0.05$.

common for antioxidant activity. These methods are also used to estimate the antioxidant activity because of the relatively short time required for analysis [34]. Many

antioxidant studies have been conducted on *L. officinalis* and reported to be related to the total amount of flavonoids and phenolic compounds. Compared with these other

studies, it has been shown to have a high antioxidant effect because of flavonoids and phenolic compounds like other studies [6, 31–34, 37].

Oxidative stress causes many diseases such as gastric diseases [38, 39]. Many studies have shown that plant components especially flavonoids can be effective and protective against oxidative damage [40, 41]. The flavonoid compositions of the fruit extract were determined by LC/MS analyses. The flavonoid components of the extract were characterized as luteolin, quercetin, and apigenin. Quercetin has been high antioxidant activity, which is a member of the flavonoid's family. It is the most effective scavenger of ROS [42]. Some studies have shown that quercetin is effective against gastric cancer [43, 44]. And also previous studies have shown that luteolin and apigenin also have effective activity for gastric cancer [45–48].

In the present study, the cytotoxic activity of the extract can be thought to be due to these flavonoid group compounds. In previous cytotoxic activity studies, it can be realized that the cytotoxicity may cause by flavonoids [49, 50]. Previous studies have shown that *L. officinalis* fruits have a selective cytotoxic effect in lung, colon, prostate, liver, and cervical cancer cell lines and also in these studies indicated that the effect was caused by the phenolic compounds and flavonoids having antioxidant effects [20–22].

Our results revealed that ranging from 5 to 10 mg/mL concentrations of the extracts induce highly significant cell death in AGS and MKN-45 cell lines whilst preserving human fibroblasts healthier. We also investigated the background of cytotoxicity of the extract in terms of cell death pattern. Regarding LC/MS findings, luteolin, quercetin and apigenin contents of *L. Officinalis* methanol extract, might lead to apoptotic cell death in gastric cancer cells when treated. Protein cleavages (PARP and caspase 3) of the apoptotic cascade showed that the apoptosis caused by the extracts follows a caspase dependent manner. Furthermore, when the apoptotic cell population was analyzed via Annexin V/PI stainings, we detected a statistically significant shift to apoptotic cell copopulation from live cell state (including both early and late apoptosis stages).

These findings indicate that as well as antioxidant activities, *L. officinalis* fruit extracts have anti-cancer effects against gastric cancer and when further studied, the active components can be an alternative or adjuvant to standard chemical drugs used in the clinics. Since many of the active phytochemicals are toxic to normal cells, cancer selective behavior of *L. officinalis* fruit extract for certain concentrations can be promising for translational approaches and *L. officinalis* can be potentially

used against several cancers upon investigation. For this reason, the results of this study have a novelty to further study in detail and suggest to examine *L. officinalis* fruit methanol extract contents as potential anti-cancer therapeutic candidates.

Research funding: None declared.

Author contributions: All authors have accepted responsibility for the entire content of this manuscript and approved its submission.

Competing interests: The authors declare no conflict of interest.

References

1. Ayla S, Okur ME, Günel MY, Özdemir EM, Çiçek Polat D, Yoltaş A, et al. Wound healing effects of methanol extract of *Laurocerasus officinalis* Roem. *Biotech Histochem* 2019;94:180–8.
2. İslam A. 'Kiraz' cherry laurel (*Prunus laurocerasus*). *N Z J Crop Hortic Sci* 2002;30:301–2.
3. Yeşilada E, Sezik E, Honda G, Takaishi Y, Takeda Y, Tanaka T. Traditional medicine in Turkey IX: folk medicine in north-west Anatolia. *J Ethnopharmacol* 1999;64:195–210.
4. Küpeli Akkol E, Kırmızıbekmez H, Küçükboyacı N, Gören AC, Yeşilada E. Isolation of active constituents from cherry laurel (*Laurocerasus officinalis* Roem.) leaves through bioassay-guided procedures. *J Ethnopharmacol* 2012;139:527–32.
5. Var M, Ayaz A. Changes in sugar composition in cherry laurel (*CV oxygemmis*) fruit during development and ripening. *Pakistan J Bot* 2004;36:389–94.
6. Kolaylı S, Küçük M, Duran C, Candan F, Dinçer B. Chemical and antioxidant properties of *Laurocerasus officinalis* Roem. (cherry laurel) fruit grown in the Black Sea region. *J Agric Food Chem* 2003;51:7489–94.
7. Colak A, Özen A, Dincer B, Güner S, Ayaz FA. Diphenolases from two cultivars of cherry laurel (*Laurocerasus officinalis* Roem.) fruits at an early stage of maturation. *Food Chem* 2005;90:801–7.
8. Livshits Z, Rao RB, Smith SW. An approach to chemotherapy-associated toxicity. *Emerg Med Clin North Am* 2014;32:167–203.
9. Demain AL, Vaishnav P. Natural products for cancer chemotherapy. *Microb Biotechnol* 2011;4:687–99.
10. Rajesh E, Sankari L, Malathi L, Krupaa J. Naturally occurring products in cancer therapy. *J Pharm BioAllied Sci* 2015;7:183.
11. Serafini M, Bellocco R, Wolk A, Ekström AM. Total antioxidant potential of fruit and vegetables and risk of gastric cancer. *Gastroenterology* 2002;123:985–91.
12. Metere A, Giacomelli L. Absorption, metabolism and protective role of fruits and vegetables polyphenols against gastric cancer. *Eur Rev Med Pharmacol Sci* 2017;21:5820–58.
13. Borek C. Dietary antioxidants and human cancer. *Integr Canc Ther* 2004;3:333–41.
14. Venerito M, Link A, Rokkas T, Malfertheiner P. Gastric cancer – clinical and epidemiological aspects. *Helicobacter* 2016; 21:39–44.
15. Suzuki H, Mori H. Gastric cancer after helicobacter pylori eradication. *Jpn J Canc Chemother* 2018;45:1123–7.

16. Guggenheim DE, Shah MA. Gastric cancer epidemiology and risk factors. *J Surg Oncol* 2013;107:230–6.
17. Ang TL, Fock KM. Clinical epidemiology of gastric cancer. *Singap Med J* 2014;55:621–8.
18. Butcher LD, den Hartog G, Ernst PB, Crowe SE. Oxidative stress resulting from *Helicobacter pylori* infection contributes to gastric carcinogenesis. *CMGH* 2017;3:316–22.
19. Díaz P, Valderrama MV, Bravo J, Quest AFG. *Helicobacter pylori* and gastric cancer: adaptive cellular mechanisms involved in disease progression. *Front Microbiol* 2018;9:1–10.
20. Aydın A, Erenler R, Yılmaz B, Tekin Ş. Antiproliferative effect of Cherry laurel. *J Turk Chem Soc Sect A Chem* 2016;3:217.
21. Demir S, Turan İ, Demir F, Ayazoglu Demir E, Aliyazicioglu Y. Cytotoxic effect of *Laurocerasus officinalis* extract on human cancer cell lines. *Marmara Pharm J* 2017;21:121–6.
22. Çakir B, Gülseren İ. Investigations on apoptotic activities of cherry laurel extracts in HCT-116 human colon carcinoma cells. *Indian J Pharm Educ Res* 2019;53:S264–72.
23. Okur ME, Ayla Ş, Çiçek-Polat D, Günel MY, Voltas A, Biçeroğlu Ö. Novel insight into wound healing properties of methanol extract of *Capparis ovata* Desf. var. *palaestina* Zohary fruits. *J Pharm Pharmacol* 2018;70:1–13.
24. Spanos GA, Wrolstad RE. Influence of processing and storage on the phenolic composition of Thompson seedless grape juice. *J Agric Food Chem* 1990;38:1565–71.
25. Singleton VL, Orthofer R, Lamuela-Raventós RM. [14] Analysis of total phenols and other oxidation substrates and antioxidants by means of folin-ciocalteu reagent. *Methods Enzymol* 1999;299:152–78.
26. Okur ME, Polat DC, Ozbek H, Yılmaz S, Voltas A, Arslan R. Evaluation of the antidiabetic property of *Capparis Ovata* Desf. Var. *Paleastina* Zoh. Extracts using in vivo and in vitro approaches. *Endocr Metab Immune Disord - Drug Targets* 2018;18:489–501.
27. Mihai CM, Mărghitaş L, Bobiş O, Dezmirean D, Tămaş M. Estimation of flavonoid content in propolis by two different colorimetric methods. *Sci Pap Anim Sci Biotechnol* 2010;43:407–10.
28. Blois MS. Antioxidant determinations by the use of a stable free radical [10]. *Nature* 1958;181:1199–200.
29. Re R, Pellegrini N, Proteggente A, Pannala A, Yang M, Rice-Evans C. Antioxidant activity applying an improved ABTS radical cation decolorization assay. *Free Radic Biol Med* 1999;26:1231–7.
30. Gulsoy-Toplan G, Goger F, Yildiz-Peko A, Gibbons S, Sariyar G, Mat A. Chemical constituents of the different parts of *Colchicum micranthum* and *C. chalconicum* and their cytotoxic activities. *Nat Prod Commun* 2018;13:535–8.
31. Orhan IE, Akkol EK. Estimation of neuroprotective effects of *Laurocerasus officinalis* Roem. (cherry laurel) by in vitro methods. *Food Res Int* 2011;44:818–22.
32. Liyana-Pathirana CM, Shahidi F, Alasalvar C. Antioxidant activity of cherry laurel fruit (*Laurocerasus officinalis* Roem.) and its concentrated juice. *Food Chem* 2006;99:121–8.
33. Karabegović IT, Stojičević SS, Veličković DT, Todorović ZB, Nikolić NČ, Lazić ML. The effect of different extraction techniques on the composition and antioxidant activity of cherry laurel (*Prunus laurocerasus*) leaf and fruit extracts. *Ind Crop Prod* 2014;54:142–8.
34. Cai Y, Luo Q, Sun M, Corke H. Antioxidant activity and phenolic compounds of 112 traditional Chinese medicinal plants associated with anticancer. *Life Sci* 2004;74:2157–84.
35. Celep E, Aydın A, Yesilada E. A comparative study on the in vitro antioxidant potentials of three edible fruits: cornelian cherry, Japanese persimmon and cherry laurel. *Food Chem Toxicol* 2012;50:3329–35.
36. Erdemoglu N, Küpeli E, Yeşilada E. Anti-inflammatory and antinociceptive activity assessment of plants used as remedy in Turkish folk medicine. *J Ethnopharmacol* 2003;89:123–9.
37. Karahalil F, Şahin H. Phenolic composition and antioxidant capacity of cherry laurel (*Laurocerasus officinalis* Roem.) sampled from Trabzon region, Turkey. *Afr J Biotechnol* 2011;10:16293–9.
38. Tandon R, Khanna RD, Dorababu M, Goel R. Oxidative stress and antioxidants status in peptic ulcer and gastric carcinoma. *Indian J Physiol Pharmacol* 2004;48:115–8.
39. Bhattacharyya A, Chattopadhyay R, Mitra S, Crowe SE. Oxidative stress: an essential factor in the pathogenesis of gastrointestinal mucosal diseases. *Physiol Rev* 2014;94:329–54.
40. Butera D, Tesoriere L, Di Gaudio F, Bongiorno A, Allegra M, Pintaudi AM, et al. Antioxidant activities of Sicilian prickly pear (*Opuntia ficus indica*) fruit extracts and reducing properties of its betalains: betanin and indicaxanthin. *J Agric Food Chem* 2002;50:6895–901.
41. Lee J-C, Kim H-R, Kim J, Jang Y-S. Antioxidant property of an ethanol extract of the stem of *Opuntia ficus-indica* var. *Saboten*. *J Agric Food Chem* 2002;50:6490–6.
42. Boots AW, Haenen GRMM, Bast A. Health effects of quercetin: from antioxidant to nutraceutical. *Eur J Pharmacol* 2008;585:325–37.
43. Borska S, Chmielewska M, Wysocka T, Drag-Zalesinska M, Zabel M, Dziegiel P, et al. In vitro effect of quercetin on human gastric carcinoma: targeting cancer cells death and MDR. *Food Chem Toxicol* 2012;50:3375–83.
44. Wang K, Liu R, Li J, Mao J, Lei Y, Wu J, et al. Quercetin induces protective autophagy in gastric cancer cells: involvement of Akt-mTOR- and hypoxia-induced factor 1 α -mediated signaling. *Autophagy* 2011;7:966–78.
45. Wu B, Zhang Q, Shen W, Zhu J. Anti-proliferative and chemosensitizing effects of luteolin on human gastric cancer AGS cell line. *Mol Cell Biochem* 2008;313:125–32.
46. Wu H, Huang M, Liu Y, Shu Y, Liu P. Luteolin induces apoptosis by up-regulating miR-34a in human gastric cancer cells. *Technol Canc Res Treat* 2015;14:747–55.
47. Shukla S, Gupta S. Apigenin: a promising molecule for cancer prevention. *Pharm Res* 2010;27:962–78.
48. Kuo C-H, Weng B-C, Wu C-C, Yang SF, Wu DC, Wang YC. Apigenin has anti-atrophic gastritis and anti-gastric cancer progression effects in *Helicobacter pylori*-infected Mongolian gerbils. *J Ethnopharmacol* 2014;151:1031–9.
49. Hirobe C, Qiao Z-S, Takeya K, Itokawa H. Cytotoxic flavonoids from *Vitex agnus-castus*. *Phytochemistry* 1997;46:521–4.
50. Seelinger G, Merfort I, Wölflle U, Schempp CM. Anti-carcinogenic effects of the flavonoid luteolin. *Molecules* 2008;13:2628–51.