

Evaluation of Patients Admitted to the Emergency Department with the Suspect of Acute Renal Colic with the Modified STONE Score

© Birdal Güllüpinar¹, © Pinar Ayvat², © Erden Erol Ünlüer¹, © Serhat Koran³

¹Clinic of Emergency, İzmir Bozyaka Training and Research Hospital, University of Health Sciences Turkey, İzmir, Turkey

²Department of Anesthesiology and Reanimation, İzmir Democracy University Faculty of Medicine, İzmir, Turkey

³Department of Family Medicine, Istanbul Medipol University Faculty of Medicine, Istanbul, Turkey

Abstract

Aim: Renal colic pain is one of the most common agonizing forms of pain that is frequently treated in emergency departments. Computed tomography (CT), which is used for the detection of kidney stones, is a costly application. Therefore, scoring systems that predict stone have been developed. This study was conducted to investigate the diagnostic accuracy of the Modified STONE Score (MSS) to predict stones.

Materials and Methods: Among those who applied to the emergency department with renal colic pain, patients with CT were examined. Three hundred and thirty-seven patients included in the study were divided into two groups as those with and without kidney stones. It was examined whether there was a difference between these two groups in terms of personal, seasonal, laboratory findings and MSS.

Results: We found that ureteral stone history, pain duration less than 6 hours, presence of hematuria and nausea/vomiting, C-reactive protein (CRP) value below 0.5 mg/dL, The MSS above 9, age ≤ 50 years were factors that increase stone. The MSS was significantly high in the stone-detected group. When the STONE score is calculated for all patients and divided into three groups (low, moderated, and high modified STONE scores), the prevalence of ureteral stones increases towards the high MSS group.

Conclusion: We found that the modified STONE score was quite successful in predicting ureteral stones. We determined that emergency physicians can diagnose stones using this score and avoid unnecessary CT. The diagnostic value of this score may increase when nausea/vomiting factor is added.

Keywords: Modified STONE Score, urinary tract stones, emergency departments

Introduction

Renal colic pain is a painful urologic case caused by the presence of stones in the urinary tract. Typically, the pain is blunt, continuous, and excruciating, with abrupt onset at the costo-vertebral angle, radiating to the groin and genitals (1). Renal colic pain affects more than 1 million patients who visit hospitals due to complaints from it each year (2) Almost half of these patients revisit hospitals within five to seven years after the first visit. More than 70% of kidney stone occurrences are observed in people aged 20 to 50 years, and the incidence in men is about two times higher than in women (3-5). In Turkey, the prevalence of ureteral stones was reported at approximately 14% (6).

Urinalysis, laboratory tests and imaging methods such as ultrasonography (USG) or computed tomography (CT) are used in renal colic. Although USG can accurately detect hydronephrosis and perinephric fluid, it has low sensitivity in showing and locating kidney stones (7). Non-contrast CT is the gold standard imaging method in the initial diagnosis of patients with ureteral stones (8). In recent years, CT with a sensitivity of 96.6% and a specificity of 94.9% has been the first-choice method as it can detect any hydroureter, hydronephrosis or ureteral edema associated with the location and size of the stone (8,9). Unlike USG, CT exposes patients to high levels of radiation and increases the long-term risk of cancer (10-13). It is estimated that there are new cancer patients due to unnecessary abdominal and pelvic CT scans (14). Therefore, various scoring systems such as STONE score, Modified



Corresponding Author: Asst. Prof. Pinar Ayvat, Department of Anesthesiology and Reanimation, İzmir Democracy University Faculty of Medicine, İzmir, Turkey
E-mail: drpinarunde@yahoo.com ORCID ID: orcid.org/0000-0002-9941-3109

Received: 29.06.2021
Accepted: 10.11.2021

Cite this article as: Güllüpinar B, Ayvat P, Ünlüer EE, Koran S. Evaluation of Patients Admitted to the Emergency Department with the Suspect of Acute Renal Colic with the Modified Stone Score. Eurasian J Emerg Med. 2022;21(2):138-45.

© Copyright 2022 by the Emergency Medicine Physicians' Association of Turkey
Eurasian Journal of Emergency Medicine published by Galenos Publishing House.

STONE Score (MSS) and CHOKAI Score have been developed to prevent unnecessary CT use and thus reduce radiation exposure in patients with ureteral stones (15-17). The STONE score has a sensitivity of 71.7% and a specificity of 64.7%, while the sensitivity and specificity of the MSS are reported as 87.7% and 70.6% (17). For this reason, many researchers preferred to use MSS.

Although there are many studies on the STONE Score in the literature, there are few studies evaluating the MSS. This study was conducted to investigate the diagnostic accuracy of the MSS and its ability to predict the presence of ureteral stones.

Materials and Methods

This study was designed as a single-center, retrospective observational study and it was carried out in İzmir Bozyaka Training and Research Hospital which has a capacity of 700 beds and an annual average of 200,000 patients. Inclusion criteria for patients were: Applied to emergency department between January 2019 and January 2020, aged 18 and older, had CT scans and diagnosed with renal colic in the emergency department. The person choosing the patient was blind to the study and just did this job. Exclusion criteria for patients were: Being younger than 18 years old, not having a CT scan, having a history of trauma, presence of active malignancy, presence of known renal disease (creatinine >1.5 mg/dL), patients with unavailable or missing laboratory data, presence of leukocytes in urine microscopy and having fever more than 37.7 °C. Demographic data (age, gender, season of application) were retrieved using the hospital information management system database. The diagnosis of patients who came to the hospital with renal colic complaints, was made by anamnesis, physical examination, laboratory findings, abdominal USG and/or abdominal tomography. The patients included in the study were divided into two groups as those with and without ureteral stones. MSS was used in patients with ureteral stones. The MSS is calculated according to the values: Gender (male-3 points), duration of pain (<6 hours-3 points), presence of hematuria (6 points), previous stone history (2 points), C-reactive protein (CRP): <5 mg/L-2 points. Afterwards, the patients were divided into three groups as low risk between 0-4 points, medium risk between 5-9 points, and high risk between 10-16 points. In addition, as in the study of Kim et al. (16), we determined the optimum cut-off value for MSS as 10. According to this cut-off value, we divided them into two groups as MSS positive and MSS-negative. The accuracy rate in the prediction of ureteral stones was examined in all groups.

Statistical Analysis

SPSS 26.0 (IBM Corp., Armonk, NY, USA) program was used in the analysis of variables. Whether the data were suitable for normal distribution was evaluated with the Shapiro-Wilk test.

Independent-samples t-test was used together with Bootstrap results to compare two independent groups with each other according to quantitative data. Mann-Whitney U test was used in conjunction with Monte Carlo results. Pearson chi-square test was tested with Monte Carlo Simulation technique to compare categorical variables with each other. Column ratios were compared with each other and expressed according to Benjamini-Hochberg corrected p value results. Odds ratio with 95% confidence intervals was used to compare patients with and without a risk factor. In order to determine the causal relationship of the presence of ureteral stone with explanatory variables, logistic regression test was tested with the Enter method. Sensitivity and specificity ratios were analyzed by receiver operating curve (ROC) curve analysis to determine the relationship between the real classification and the classification calculated with the cut off values according to the MSS, age and CRP variables. Quantitative variables were shown in the tables as mean (standard deviation) and median (percentile 25/percentile 75), and categorical variables as n (%). Variables were analyzed at 95% confidence level, and p value less than 0.05 was considered statistically significant.

Results

Of 1165 patients admitted to the emergency department with suspected acute renal colic, 337 met the inclusion criteria (Figure 1).




Figure 1. Flow diagram of the study
CR: Creatinine, CT: Computed tomography

In this study, the median (percentile 25/percentile 75) age was 41 (33/52) and the number of male patients was 226 (67.1%). Pain duration was less than six hours in 59.3% of the patients, and it was accompanied by nausea and/or vomiting in 55.5%. Those with a history of ureteral stone in the past were 41.2%. The most common laboratory finding was hematuria with 70.0%. According to CT scan findings, ureteral stones were detected in 237 (70.3%) patients. History of ureteral stone, pain duration less than 6 hours, presence of hematuria, CRP value below 0.5 mg/dL, and STONE score above 9 were significantly more common in the group with ureteral stones ($p < 0.001$). In addition, a statistically significant difference was found in the stone-detected group compared to the stone-free group in terms of age (≤ 50) and the presence of nausea and/or vomiting ($p = 0.011$, $p = 0.008$). While the median (minimum/maximum) value of the MSS was 12 (11/14) in the stone-detected group, it was 4.5 (3/6) in the other group, and the difference between the groups was statistically significant ($p < 0.001$). No significant difference was found between the groups in terms of leukocyte count (< 12.000 cells/mL), BUN (< 26 mg/dL) value and creatinine (< 1.2 mg/dL)

value, the time for application to hospital and gender ($p > 0.05$) (Table 1), (Figure 2). Logistic regression test revealed that 3 factors were statistically significant ($p < 0.001$): Age ≤ 50 [odds ratio (OR)=5.542], presence of nausea and vomiting (OR=6.83), MSS > 9 (OR=100.048) were associated with the incidence of renal stone (Table 2).

According to the MSS, 233 (69.1%) patients were in the high-risk group, 56 (16.6%) patients in the moderated risk group, and 48 (14.2%) patients in the low-risk group. The prevalence of ureteral stones was 227/233 (97.4%) in the high-risk group, 10/56 (17.8%) in the moderated-risk group, and 0/48 (0%) in the low-risk group. As we move from the low-risk group to the high-risk group, the prevalence of ureteral stones increases (Figure 3).

When the MSS cut-off value was accepted as 10, those with MSS 9 and below were called MSS-negative, those with MSS 10 and above were called MSS-positive. The relationship between the diagnosis of renal stones and the diagnosis of alternative diseases between these two groups is shown in Figure 4.




Figure 2. ROC curves about Modified STONE Score, age and CRP level of the patients

ROC: Receiver operating characteristics, CRP: C-reactive protein




Figure 3. Comparison of patients with low, moderated and high modified STONE Scores in terms of detection of ureteral stones and alternative diagnoses




Figure 4. Comparison of patients with positive and negative modified STONE Scores in terms of detecting ureteral stones and alternative diagnoses

MSS: Modified STONE Score

Table 1. Comparison of demographic data, laboratory findings, Modified STONE Score findings of patients with and without ureteral stone

| | Total (n=337) | Patients without stone (n=100) | Patients with stone (n=237) | p value |
|------------------------------------|--------------------|-----------------------------------|--------------------------------|---------------------------------|
| Age, median (q1/q3) | 41 (33/52) | 43 (35/56) | 40 (32/50) | 0.011^u |
| Gender, n (%) | | | | |
| Female | 111 (32.9) | 40 (40.0) | 71 (30.0) | 0.077 ^c |
| Male | 226 (67.1) | 60 (60.0) | 166 (70.0) | |
| Season, n (%) | | | | |
| Autumn | 73 (21.7) | 21 (21.0) | 52 (21.9) | 0.354 ^c |
| Winter | 87 (25.8) | 30 (30.0) | 57 (24.1) | |
| Spring | 69 (20.5) | 15 (15.0) | 54 (22.8) | |
| Summer | 108 (32.0) | 34 (34.0) | 74 (31.2) | |
| Pain duration, n (%) | | | | |
| <6 | 200 (59.3) | 25 (25.0) | 175 (73.8) ^A | 8.5 (4.9-14.5) ^{OR} |
| >6 | 137 (40.7) | 75 (75.0) ^B | 62 (26.2) | |
| Nausea, vomiting, n (%) | | | | |
| No | 150 (44.5) | 56 (56.0) ^B | 94 (39.7) | 1.9 (1.2-3.1) ^{OR} |
| Yes | 187 (55.5) | 44 (44.0) | 143 (60.3) ^A | |
| Stone story, n (%) | | | | |
| No | 198 (58.8) | 82 (82.0) ^B | 116 (48.9) | 4.7 (2.7-8.4) ^{OR} |
| Yes | 139 (41.2) | 18 (18.0) | 121 (51.1) ^A | |
| Hematuria, n (%) | | | | |
| No | 101 (30.0) | 83 (83.0) ^B | 18 (7.6) | 59.4 (29.2-120.7) ^{OR} |
| Yes | 236 (70.0) | 17 (17.0) | 219 (92.4) ^A | |
| Creatinine, mean (SD) | 1.1 (0.2) | 1 (0.2) | 1.1 (0.2) | 0.485 ^t |
| BUN, median (q1/q3) | 31 (25/37) | 30.5 (24/40) | 31 (25/37) | 0.893 ^u |
| Urea, median (q1/q3) | 66.34 (53.5/79.18) | 65.27 (51.36/85.6) | 66.34 (53.5/79.18) | 0.893 ^u |
| CRP, median (q1/q3) | 0.4 (0.2/3.7) | 1.6 (0.3/7.25) | 0.4 (0.2/2.8) | <0.001^u |
| WBC, median (q1/q3) | 9.47 (7.59/12.07) | 9.775 (7.57/13.22) | 9.43 (7.61/11.39) | 0.155 ^u |
| STONE Score, median (q1/q3) | 11 (7/13) | 4.5 (3/6) | 12 (11/14) | <0.001^u |
| Age, n (%) | | | | |
| >50 | 91 (27.0) | 37 (37.0) ^{sp} | 54 (22.8) | AUC (SE): 0.588 (0.034) |
| ≤50 | 246 (73.0) | 63 (63.0) | 183 (77.2) ^{ss} | |
| CRP, n (%) | | | | |
| >0.4 | 146 (43.3) | 63 (63.0) ^{sp} | 83 (35.0) | AUC (SE): 0.668 (0.031) |
| ≤0.4 | 191 (56.7) | 37 (37.0) | 154 (65.0) ^{ss} | |
| STONE Score, n (%) | | | | |
| ≤9 | 104 (30.9) | 94 (94.0) ^{sp} | 10 (4.2) | AUC (SE): 0.971 (0.011) |
| >9 | 233 (69.1) | 6 (6.0) | 227 (95.8) ^{ss} | |

^uMann Whitney U test (Monte Carlo), ^cPearson chi-square test (Monte Carlo), ^{OR}Odds Ratio (%95 confidence interval), ^tReceiver operating characteristics (ROC) curve analysis (Honley & McNell-Youden index J), AUC: Area under the ROC curve, q1: percentile 25, q3: percentile 75, ^ASignificance according to the "patients with stones" group, ^BSignificance according to the "patients without stones" group, Significant values are shown in bold.
BUN: Blood urea nitrogen, CRP: C-reactive protein, WBC: White blood cells, SD: Standard deviation, SE: Standard error, n: Number

Table 2. Odds ratio for age, presence of nausea and vomiting, and Modified STONE Score

| | B | SE | p value | Odds ratio | Odds ratio (95% CI) | |
|----------------------|--------|-------|---------|------------|---------------------|---------|
| | | | | | Lower | Upper |
| Age (≥50) | 1.712 | 0.459 | <0.001 | 5.542 | 2.253 | 13.631 |
| Nausea, vomiting | 1.921 | 0.341 | <0.001 | 6.830 | 3.504 | 13.315 |
| Modified STONE Score | -4.606 | 0.547 | <0.001 | 100.048 | 34.230 | 292.418 |

Dependent variable: presence of stones, prediction rate of patients with stones=96.2, predictions of patients with non-stones=83, overall accuracy: 92.3, p model: <0.001. Multiple Logistic Regression (Method=Enter), CI: Confidence interval, B: Regression coefficients, SE: Standard error

Alternative diagnoses according to ureteral stone prevalence and MSS categories are shown in Table 3.

While the imaging results were completely normal in 59 (17.5%) of 100 (29.7%) patients who do not have ureteral stones according to CT scans, pathologies other than ureteral stones were detected in 41 (12.2%) patients. Other pathologies include alternative diagnoses such as acute appendicitis, acute cholecystitis, acute pancreatitis, ovarian cyst rupture, renal infarction, gastrointestinal perforation, etc. (Table 4).

Discussion

This study demonstrated the applicability of the MSS in the Turkish population in patients admitted to the emergency department with suspected acute renal colic. Ureteral stones were detected at a rate of 97.4% in the group with a high MSS. This result was similar with the rate of 98% found in the study of Kim et al. (16).

History and physical examination findings are very important in patients presenting to the emergency department, but emergency physicians can use scoring systems as a complementary tool. It was showed in our study that the MSS had a high sensitivity in detecting ureteral stones. Patients with a high MSS are more likely to have ureteral stones. For this reason, patients can be diagnosed without the CT, USG or extra consultation. Thus, rapid discharge of the patient from the emergency department can be planned. For this reason, we think that this scoring system can be preferred by emergency physicians and will reduce the workload and additional cost.

In our retrospective study, 1165 patients with suspected acute renal colic were screened and 598 patients had CT scans. CT imaging method was used in 51.3% of the patients in our own clinic. The rate of CT imaging in our clinic was similar to the rate found in the study of Kim et al. (16).

In our study, the prevalence of ureteral stones was 70.3% whereas in the study conducted by Kim et al. (16) in South Korea, the prevalence of ureteral stones was 79%. In Turkey, the prevalence of ureteral stones varies between 49-84% (17-19).

Table 3. Alternative diagnoses in patients according to modified STONE score categories

| | Low risk (n=19) | Moderate risk (n=18) | High risk (n=4) | Total (n=41) |
|-----------------------------|-----------------|----------------------|-----------------|--------------|
| Acute appendicitis | 1 | - | - | 1 |
| Cholelithiasis | 2 | 2 | - | 4 |
| Newly detected malignancy | 2 | 1 | - | 3 |
| Acute pyelonephritis | 2 | 2 | - | 4 |
| Acute cholecystitis | 1 | 1 | - | 2 |
| Ovarian cyst | 1 | 2 | - | 3 |
| Enterocolitis | 1 | - | 1 | 2 |
| Inguinal hernia | - | 2 | 1 | 3 |
| Ureteropelvic stenosis | - | 1 | - | 1 |
| Renal infarction | - | 1 | - | 1 |
| Ovarian cyst rupture | 1 | - | - | 1 |
| Pelvic inflammatory disease | - | - | 1 | 1 |
| GIS perforation | 1 | 1 | - | 2 |
| Endometriosis | - | 1 | - | 1 |
| Thoracolumbar spondylosis | 1 | - | - | 1 |
| Acute pancreatitis | - | 1 | - | 2 |
| Angiomyolipoma | 1 | - | - | 1 |
| Paraganglioma | 1 | - | - | 1 |
| Hydatid cyst of the liver | 1 | - | - | 1 |
| Appendicolith | - | 1 | - | 1 |
| Diverticulitis | - | 1 | 1 | 2 |
| Adrenal adenoma | 1 | 1 | - | 2 |
| Aortic syndrome | 1 | - | - | 1 |
| Pneumonia | 1 | - | - | 1 |

n: Number

The MSS included the variables male sex, pain duration less than six hours, previous ureter stone history, presence of hematuria, and CRP less than 0.5 mg/dL (16). In our study, all these parameters were the same as the other study findings except gender (16,17). Among the patients included in our study, 226 patients (67.1%) were male. While male gender was an important risk factor compared to females in other studies (16,17), although a high rate of male patients was included in our study, the difference was statistically not significant ($p=0.07$). Since there is no obstetrics and gynecology department in our hospital, female patients in our region often apply to the emergency departments of other hospitals in the region when they are ill. Therefore, the number of male patients could be higher in our hospital.

Although male gender was a risk factor according to the MSS, it was not a risk factor in our study. This may be a difference seen in our society.

In our study, the prevalence of ureteral stones was 227/233 (97.4%) in the high-risk group, 10/56 (17.8%) in the intermediate-risk group, and 0/48 (0%) in the low-risk group. The prevalence of alternative diagnosis is increasing from the high-risk group to the low- and intermediate-risk group. In the alternative diagnosis group, the prevalence was found to be 39.6% (19/48) for the low-risk group, 32.1% (18/56) for the medium risk group, and 1.7% (4/233) for the high-risk group. These findings also showed similarities as in the study of Kim et al. (16).

Table 4. Alternative diagnoses in patients without ureteral stones

| | n | % |
|--|-----|------|
| Patients having stone | 237 | 70.3 |
| Patients not having stone and not being detected any diseases | 59 | 17.5 |
| Patients not having stone but being detected alternative diseases | 41 | 12.2 |
| Cholelithiasis | 4 | 1.2 |
| Pyelonephritis | 4 | 1.2 |
| Inguinal hernia | 3 | 0.9 |
| Ovarian cyst | 3 | 0.9 |
| New malignancy | 3 | 0.9 |
| Diverticulitis | 2 | 0.6 |
| Enterocolitis | 2 | 0.6 |
| Gastrointestinal perforation | 2 | 0.6 |
| Cholecystitis | 2 | 0.6 |
| Adrenal adenoma | 2 | 0.6 |
| Angiomyolipoma | 1 | 0.3 |
| Aortic syndrome | 1 | 0.3 |
| Appendicitis | 1 | 0.3 |
| Appendicolith | 1 | 0.3 |
| Endometriosis | 1 | 0.3 |
| Hydatid cyst of the liver | 1 | 0.3 |
| Ovarian cyst rupture | 1 | 0.3 |
| Pancreatitis | 1 | 0.3 |
| Paraganglioma | 1 | 0.3 |
| Pelvic inflammatory disease | 1 | 0.3 |
| Pneumonia | 1 | 0.3 |
| Renal infarction | 1 | 0.3 |
| Thoracolumbar spondylosis | 1 | 0.3 |
| Ureteropelvic stenosis | 1 | 0.3 |

We found the alternative diagnosis rate to be 12.1% in patients who applied to our emergency department with the suspicion of acute renal colic. In similar studies, the rate of alternative diagnosis varies between 10% and 22.1% (20-22). The higher rate of alternative diagnosis in other countries compared to our country can be explained by the more widespread use of CT imaging.

As in the study of Kim et al. (16), we determined the optimum cut-off value for MSS as 10, and according to this cut-off value, two groups were designated as MSS positive and MSS-negative. The MSS positive group corresponds to the high-risk group and includes patients with a MSS of 10 or more. The prevalence of ureteral stones in the MSS positive group was 97.4% in our study, while it was 98.0% in the study of Kim et al. (16).

In our study, important alternative diagnoses were found to be 1.7% in the MSS positive group. These diagnoses were diverticulitis, inguinal hernia, enterocolitis and pelvic inflammatory disease. In the study of Kim et al. (16), the rate of significant alternative diagnosis in the MSS positive group was 1.9% which was similar to the result in our study. On the other hand, the rate of significant alternative diagnosis in the MSS negative group was found to be 35.6% in our study. In the study of Kim et al. (16), this rate was 23.5%. Therefore, it can be said that advanced imaging methods are necessary for patients in the MSS negative group.

It is known that acute renal colic pain is more common in young adult men and often recurs (3-5). Therefore, young patients have frequent admissions to the emergency department. There are studies which showed that patients are exposed to high levels of radiation with CT and this increased the risk of cancer, particularly in younger patients compared to the elderly (23). In addition, unnecessary IT requests both increase the length of stay in the emergency department and

cause additional costs. As seen in our study and other studies, MSS positivity largely detects ureteral stones, and it can be said that additional imaging methods other than ultrasound can be abandoned.

In our study, we detected ureteral stones in 67.4% (233/347) of the patients with MSS positive. In the study of Kim et al. (16), this rate was 64.2%. Accordingly, it can be interpreted that using MSS can reduce the use of CT method by 60-70%, and thus reduce the risk of cancer due to radiation.

Study Limitations

Our study has some limitations. The sample size was relatively small and it was a retrospective study. In addition, the generalizability of the findings is limited as it is a single-center study. The results may differ from region to region or country to country due to differences such as geographical conditions, socioeconomic level, eating and drinking habits that may affect the symptom of acute renal colic. Our study was only for a part of the Turkish population. The results of this study should be supported by multicenter studies that can cover all regions of the country or studies covering different countries. Only patients who underwent CT imaging in the MSS evaluation were examined. However, other imaging methods are also frequently used in our country. These limitations should be considered in future research.

Conclusion

With this study, we found that the MSS is quite successful in predicting ureteral stones. By using this score, emergency physicians will safely reduce the overuse of CT, the costs, and the length of stay in the emergency room. We think that when factors such as nausea and/or vomiting are added to this score, its diagnostic value may increase. The sensitivity and specificity of the MSS should be supported by more extensive studies.

Ethics

Ethics Committee Approval: This study was approved by İstanbul Medipol University Noninvasive Clinical Researches Ethics Committee (decision no: 440, date: 15.04.2021).

Informed Consent: Retrospective study.

Peer-review: Externally peer-reviewed.

Authorship Contributions

Surgical and/or Medical Practices: B.G., E.E.Ü., Concept: E.E.Ü., Design: B.G., P.A., E.E.Ü., S.K., Data Collection and/or Processing: B.G., P.A., S.K., Analysis and/or Interpretation: B.G., P.A., E.E.Ü., Literature Search: B.G., P.A., E.E.Ü., S.K., Writing: B.G., P.A., S.K.

Conflict of Interest: No conflict of interest was declared by the authors.

Financial Disclosure: The authors declared that this study received no financial support.

References

- Bultitude M, Rees J. Management of renal colic. *BMJ*. 2012;345:e5499.
- Golzari SE, Soleimanpour H, Rahmani F, Zamani Mehr N, Safari S, Heshmat Y, et al. Therapeutic approaches for renal colic in the emergency department: a review article. *Anesthesiol Pain Med*. 2014;4:e16222.
- Agenosov MP, Kagan OF, Kheifets VK. Features of urolithiasis in patients of advanced and senile age. *Adv Gerontol*. 2018;31:368-73.
- D'Costa MR, Pais VM, Rule AD. Leave no stone unturned: defining recurrence in kidney stone formers. *Curr Opin Nephrol Hypertens*. 2019;28:148-53.
- Raja AS, Pourjabbar S, Ip IK, Baugh CW, Sodickson AD, O'Leary M, et al. Impact of a Health Information Technology-Enabled Appropriate Use Criterion on Utilization of Emergency Department CT for Renal Colic. *AJR Am J Roentgenol*. 2019;212:142-5.
- Romero V, Akpınar H, Assimos DG. Kidney stones: a global picture of prevalence, incidence, and associated risk factors. *Rev Urol* 2010; 12: pp. e86-96.
- Watkins S, Bowra J, Sharma P, Holdgate A, Giles A, Campbell L. Validation of emergency physician ultrasound in diagnosing hydronephrosis in ureteric colic. *Emerg Med Australas*. 2007;19:188-95.
- Turk C, Petrik A, Sarica K, Seitz C, Skolarikos A, Straub M, et al. EAU Guidelines on Urolithiasis. *Eur Urol*. 2016;69:475-82.
- Brisbane W, Bailey MR, Sorensen MD. An overview of kidney stone imaging techniques. *Nat Rev Urol*. 2016;13:654-62.
- National Research Council Health risks from exposure to low levels of ionizing radiation: BEIR VII phase 2. Washington, DC: National Academies Press, 2006.
- Preston DL, Ron E, Tokuoka S, Funamoto S, Nishi N, Soda M, et al. Solid cancer incidence in atomic bomb survivors: 1958–1998. *Radiat Res*. 2007;168:1-64.
- Pearce MS, Salotti JA, Little MP, McHugh K, Lee C, Kim KP, et al. Radiation exposure from CT scans in childhood and subsequent risk of leukaemia and brain tumours: a retrospective cohort study. *Lancet*. 2012;380:499-505.
- Mathews JD, Forsythe AV, Brady Z, Butler MW, Goergen SK, Byrnes GB, et al. Cancer risk in 680,000 people exposed to computed tomography scans in childhood or adolescence: data linkage study of 11 million Australians. *BMJ*. 2013;346:f2360.
- Berrington de González A, Mahesh M, Kim KP, Bhargavan M, Lewis R, Mettler F, et al. Projected cancer risks from computed tomographic scans performed in the United States in 2007. *Arch Intern Med*. 2009;169:2071-7.
- Moore CL, Bomann S, Daniels B, Luty S, Molinaro A, Singh D, et al. Derivation and validation of a clinical prediction rule for uncomplicated ureteral STONE--the STONE score: retrospective and prospective observational cohort studies. *BMJ*. 2014; 348: pp. g2191.
- Kim B, Kim K, Kim J, Jo YH, Lee JH, Hwang JE, et al. External validation of the STONE score and derivation of the modified STONE score. *Am J Emerg Med*. 2016;34:1567-72.
- Acar YA, Uysal E. External validation of STONE, modified STONE, and CHOKAI scores for the diagnosis of ureteral stones in the Turkish population. *Hong Kong J Emerg Med*. 2020; doi: 10.1177/1024907920945476.
- Rohat A, Kurt E, Şenel Ç. The comparison of two prediction models for ureteral stones: CHOKAI and STONE scores. *Am J Emerg Med*. 2020. doi: 10.1016/j.ajem.2020.08.099 S0735675720307932
- Uzun A, Korkut M, Kartal M, Bedel C. Evaluation of modified STONE score in patients presenting to the emergency department with flank pain. *Urol Sci*. 2020;31:5.

20. Moore CL, Daniels B, Singh D, Luty S, Molinaro A. Prevalence and clinical importance of alternative causes of symptoms using a renal colic computed tomography protocol in patients with flank or back pain and absence of pyuria. *Acad Emerg Med.* 2013;20:470-8.
21. Samim M, Goss S, Luty S, Weinreb J, Moore C. Incidental findings on CT for suspected renal colic in emergency department patients: prevalence and types in 5,383 consecutive examinations. *J Am Coll Radiol.* 2015;12:63-9.
22. Lokken RP, Sadow CA, Silverman SGS. Diagnostic yield of CT urography in the evaluation of young adults with hematuria. *AJR Am J Roentgenol.* 2012;198:609-15.
23. Sodickson A, Baeyens PF, Andriole KP, Prevedello LM, Nawfel RD, Hanson R, et al. Recurrent CT, cumulative radiation exposure, and associated radiation-induced cancer risks from CT of adults. *Radiology.* 2009;251:175-84.